

Rabbit PEA15 Polyclonal Antibody | anti-PEA15 antibody

Rabbit anti-PEA15 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA222722	Rabbit	Human, Mouse, Rat	Immunoprecipitation, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

PEA15; Polyclonal Antibody; Rabbit anti-PEA15 Antibody; Astrocytic phosphoprotein PEA-15; 15 kDa phosphoprotein enriched in astrocytes; Phosphoprotein enriched in diabetes; PED; anti-PEA15 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-PEA15 antibody	IP (Immunoprecipitation), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human PEA15. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of PEA15 staining in MDA-MB-231 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of PEA15 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of PEA15 expression in MDA-MB231 (A), human kidney (B), NIH3T3 (C) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-PEA15 antibody	Primary antibodies
---	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	4505705
NCBI GeneID	8682
NCBI Accession #	NP_003759.1
NCBI GenBank Nucleotide #	NM_003768.3
UniProt Accession #	Q15121; Q62048; Q5U318
Molecular Weight	15,040 Da
NCBI Official Full Name	astrocytic phosphoprotein PEA-15
NCBI Official Synonym Full Names	phosphoprotein enriched in astrocytes 15
NCBI Official Symbol	PEA15
NCBI Official Synonym Symbols	PED; MAT1; HMAT1; MAT1H; PEA-15; HUMMAT1H
NCBI Protein Information	astrocytic phosphoprotein PEA-15; homolog of mouse MAT-1 oncogene; phosphoprotein enriched in diabetes; Phosphoprotein enriched in astrocytes, 15kD; 15 kDa phosphoprotein enriched in astrocytes
UniProt Protein Name	Astrocytic phosphoprotein PEA-15
UniProt Gene Name	PEA15
UniProt Synonym Gene Names	PED
UniProt Entry Name	PEA15_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/pea15-antibody-immunoprecipitation-immunofluorescence-immunohistochemistry-western-blot-elisa/222722>