

Rabbit CBFB Polyclonal Antibody | anti-CBFB antibody

Rabbit anti-CBFB Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA222744	Rabbit	Human, Pig, Zebrafish	Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

CBFB; Polyclonal Antibody; Rabbit anti-CBFB Antibody; Core-binding factor subunit beta; CBF-beta; Polyomavirus enhancer-binding protein 2 beta subunit; PEA2-beta; PEBP2-beta; SL3-3 enhancer factor 1 subunit beta; SL3/AKV core-binding factor beta subunit; anti-CBFB antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Pig, Zebrafish
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-CBFB antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human CBFB. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of CBFB staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of CBFB staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of CBFB expression in Jurkat (A), HUVEC (B), K562 (C) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-CBFB antibody	Primary antibodies
--	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	13124873
NCBI GeneID	865
NCBI Accession #	NP_001746.1
NCBI GenBank Nucleotide #	NM_001755.2
UniProt Accession #	Q13951; Q13124; Q9HCT2; A8K347
Molecular Weight	21,508 Da
NCBI Official Full Name	core-binding factor subunit beta isoform 2
NCBI Official Synonym Full Names	core-binding factor, beta subunit
NCBI Official Symbol	CBFB
NCBI Official Synonym Symbols	PEBP2B
NCBI Protein Information	core-binding factor subunit beta; CBF-beta; PEA2-beta; PEBP2-beta; SL3-3 enhancer factor 1 beta subunit; SL3-3 enhancer factor 1 subunit beta; SL3/AKV core-binding factor beta subunit; polyomavirus enhancer-binding protein 2 beta subunit; polyomavirus enh
UniProt Protein Name	Core-binding factor subunit beta
UniProt Gene Name	CBFB
UniProt Synonym Gene Names	CBF-beta; PEA2-beta; PEBP2-beta
UniProt Entry Name	PEBB_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://stage.aaabiotech.com/polyclonal/cbfb-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/222744>