

Rabbit HNF4 alpha/gamma Polyclonal Antibody | anti-HNF4alpha/gamma antibody

Rabbit anti-HNF4 alpha/gamma Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA222808	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

HNF4 alpha/gamma; Polyclonal Antibody; Rabbit anti-HNF4 alpha/gamma Antibody; HNF4A; HNF4; NR2A1; TCF14; Hepatocyte nuclear factor 4-alpha; HNF-4-alpha; Nuclear receptor subfamily 2 group A member 1; Transcription factor 14; TCF-14; Transcription factor HNF-4; HNF4G; NR2A2; Hepatocyte nuclear factor 4-gamma; HNF-4-gamma; Nuclear receptor subfamily 2 group A member 2; anti-HNF4alpha/gamma antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-HNF4alpha/gamma antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human HNF4 alpha/gamma. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES



IF (Immunofluorescence)

Immunofluorescent analysis of HNF4 alpha/gamma staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of HNF4 alpha/gamma staining in human colon cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of HNF4 alpha/gamma expression in A549 (A), HepG2 (B), HeLa (C) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-HNF4alpha/gamma antibody	Primary antibodies
---	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	31077205
NCBI GeneID	3172
NCBI Accession #	NP_000448.3
NCBI GenBank Nucleotide #	NM_000457.4
UniProt Accession #	P41235; Q14541; P49698; Q9WUU6; P22449
Molecular Weight	52,785 Da
NCBI Official Full Name	hepatocyte nuclear factor 4-alpha isoform b
NCBI Official Synonym Full Names	hepatocyte nuclear factor 4, alpha
NCBI Official Symbol	HNF4A
NCBI Official Synonym Symbols	TCF; HNF4; MODY; MODY1; NR2A1; TCF14; HNF4a7; HNF4a8; HNF4a9; NR2A21; HNF4alpha
NCBI Protein Information	hepatocyte nuclear factor 4-alpha; TCF-14; HNF4alpha10/11/12; transcription factor 14; transcription factor HNF-4; hepatic nuclear factor 4 alpha; nuclear receptor subfamily 2 group A member 1
UniProt Protein Name	Hepatocyte nuclear factor 4-alpha
UniProt Gene Name	HNF4A
UniProt Synonym Gene Names	HNF4; NR2A1; TCF14; HNF-4-alpha; TCF-14
UniProt Entry Name	HNF4A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/hnf4-alpha-gamma-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/222808>