

Rabbit HSD17B2 Polyclonal Antibody | anti-HSD17B2 antibody

Rabbit anti-HSD17B2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA223014	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

HSD17B2; Polyclonal Antibody; Rabbit anti-HSD17B2 Antibody; EDH17B2; Estradiol 17-beta-dehydrogenase 2; 17-beta-hydroxysteroid dehydrogenase type 2; 17-beta-HSD 2; 20 alpha-hydroxysteroid dehydrogenase; 20-alpha-HSD; E2DH; Microsomal 17-beta-hydroxysteroid dehydrogenase; Testosterone 17-beta-dehydrogenase; anti-HSD17B2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-HSD17B2 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant full length protein of human HSD17B2
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of HSD17B2 staining in SW480 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of HSD17B2 staining in rat brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of HSD17B2 expression in SW480 (A), mouse liver (B), mouse kidney (C) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-HSD17B2 antibody

Primary antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	4504503
NCBI GeneID	3294
NCBI Accession #	NP_002144.1
NCBI GenBank Nucleotide #	NM_002153.2
UniProt Accession #	P37059; P51658; Q62730
Molecular Weight	42,785 Da
NCBI Official Full Name	estradiol 17-beta-dehydrogenase 2
NCBI Official Synonym Full Names	hydroxysteroid (17-beta) dehydrogenase 2
NCBI Official Symbol	HSD17B2
NCBI Official Synonym Symbols	HSD17; SDR9C2; EDH17B2
NCBI Protein Information	estradiol 17-beta-dehydrogenase 2; E2DH; 20-alpha-HSD; 17-beta-HSD 2; testosterone 17-beta-dehydrogenase; 20 alpha-hydroxysteroid dehydrogenase; 17-beta-hydroxysteroid dehydrogenase type 2; microsomal 17-beta-hydroxysteroid dehydrogenase; short chain dehy
UniProt Protein Name	Estradiol 17-beta-dehydrogenase 2
UniProt Gene Name	HSD17B2
UniProt Synonym Gene Names	EDH17B2; 17-beta-HSD 2
UniProt Entry Name	DHB2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/hsd17b2-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/223014>