

Rabbit Bcl-2 Polyclonal Antibody | anti-Bcl-2 antibody

Bcl-2 Polyclonal Antibody

CATALOG # AAA22304	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Western Blot, Immunofluorescence
------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

Bcl-2; Polyclonal Antibody; Bcl-2 Polyclonal Antibody; PPP1R50; BCL2; anti-Bcl-2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.01% thiomersal, 50% glycerol, pH 7.3.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-Bcl-2 antibody	WB (Western Blot), IF (Immunofluorescence)
Application Notes	WB: 1:500-1:2000 IF: 1:50-1:200
Immunogen	Recombinant fusion protein of human Bcl-2
Research Area	Cancer, Cell Biology, Metabolism, Signal Transduction
Preparation and Storage	Storage: -20 degree C/One year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescence analysis of A-549 cells using Bcl-2 Polyclonal Antibody at dilution of 1:200. Blue: DAPI for nuclear staining.




IF (Immunofluorescence)

Immunofluorescence analysis of U2OS cells using Bcl-2 Polyclonal Antibody at dilution of 1:200. Blue: DAPI for nuclear staining.




IF (Immunofluorescence)

Immunofluorescence analysis of C6 cells using Bcl-2 Polyclonal Antibody at dilution of 1:200. Blue: DAPI for nuclear staining.




IF (Immunofluorescence)

Immunofluorescence analysis of HeLa cells using Bcl-2 Polyclonal Antibody at dilution of 1:200. Blue: DAPI for nuclear staining.



IF (Immunofluorescence)

Immunofluorescence analysis of NIH/3T3 cells using Bcl-2 Polyclonal Antibody at dilution of 1:200. Blue: DAPI for nuclear staining.



WB (Western Blot)

Western blot analysis of extracts of various cell lines using Bcl-2 Polyclonal Antibody at 1:1000 dilution.

ADDITIONAL INFORMATION

Related Product Information for anti-Bcl-2 antibody	This gene encodes an integral outer mitochondrial membrane protein that blocks the apoptotic death of some cells such as lymphocytes. Constitutive expression of BCL2, such as in the case of translocation of BCL2 to Ig heavy chain locus, is thought to be the cause of follicular lymphoma. Alternative splicing results in multiple transcript variants.
Product Categories/Family for anti-Bcl-2 antibody	Antibodies ; Primary Antibodies ; Polyclonal Antibody

NCBI AND UNIPROT INFORMATION

NCBI GI #	72198189
NCBI GeneID	596
NCBI Accession #	NP_000624.2
NCBI GenBank Nucleotide #	NM_000633.2
UniProt Accession #	P10415 ; P10416 ; Q13842 ; Q16197
Molecular Weight	26,266 Da
NCBI Official Full Name	apoptosis regulator Bcl-2 alpha isoform
NCBI Official Synonym Full Names	B-cell CLL/lymphoma 2
NCBI Official Symbol	BCL2

NCBI Official Synonym Symbols	Bcl-2; PPP1R50
NCBI Protein Information	apoptosis regulator Bcl-2; protein phosphatase 1, regulatory subunit 50
UniProt Protein Name	Apoptosis regulator Bcl-2
UniProt Gene Name	BCL2
UniProt Entry Name	BCL2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/bcl-2-antibody-western-blot-immunofluorescence-elisa/22304>