

Rabbit RIPK2 Polyclonal Antibody | anti-RIPK2 antibody

Rabbit anti-RIPK2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA223049	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

RIPK2; Polyclonal Antibody; Rabbit anti-RIPK2 Antibody; CARDIAK; RICK; RIP2; Receptor-interacting serine/threonine-protein kinase 2; CARD-containing interleukin-1 beta-converting enzyme-associated kinase; CARD-containing IL-1 beta ICE-kinase; RIP-like-interacting CLARP kinase; Receptor-interacting protein 2; RIP-2; Tyrosine-protein kinase RIPK2; anti-RIPK2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-RIPK2 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant full length protein of human RIPK2
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of RIPK2 staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of RIPK2 staining in rat kidney formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of RIPK2 expression in MCF7 (A), mouse spleen (B), mouse testis (C) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-RIPK2 antibody

Primary antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	4506537
NCBI GeneID	8767
NCBI Accession #	NP_003812.1
NCBI GenBank Nucleotide #	NM_003821.5
UniProt Accession #	O43353; P58801
Molecular Weight	61,195 Da
NCBI Official Full Name	receptor-interacting serine/threonine-protein kinase 2
NCBI Official Synonym Full Names	receptor-interacting serine-threonine kinase 2
NCBI Official Symbol	RIPK2
NCBI Official Synonym Symbols	CCK; RICK; RIP2; CARD3; GIG30; CARDIAK
NCBI Protein Information	receptor-interacting serine/threonine-protein kinase 2; RIP-2; CARD-carrying kinase; growth-inhibiting gene 30; tyrosine-protein kinase RIPK2; receptor-interacting protein 2; CARD-containing IL-1 beta ICE-kinase; CARD-containing interleukin-1 beta-convert
UniProt Protein Name	Receptor-interacting serine/threonine-protein kinase 2
UniProt Gene Name	RIPK2
UniProt Synonym Gene Names	CARDIAK; RICK; RIP2; CARD-containing IL-1 beta ICE-kinase; RIP-2
UniProt Entry Name	RIPK2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/ripk2-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/223049>