

Rabbit BRG1 Polyclonal Antibody | anti-BRG1 antibody

Rabbit anti-BRG1 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA223057	Rabbit	Human, Mouse, Rat	Immunoprecipitation, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

BRG1; Polyclonal Antibody; Rabbit anti-BRG1 Antibody; BAF190A; SNF2B; SNF2L4; Transcription activator BRG1; ATP-dependent helicase SMARCA4; BRG1-associated factor 190A; Mitotic growth and transcription activator; Protein BRG-1; Protein brahma homolog 1; SNF2-beta; SWI/SNF-related matrix-associated actin-dependent regulator of chromatin subfamily A member 4; anti-BRG1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-BRG1 antibody	IP (Immunoprecipitation), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide of human BRG1
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of BRG1 staining in Hela cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of BRG1 staining in human colon cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of BRG1 expression in HeLa (A), K562 (B), NIH3T3 (C) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-BRG1 antibody

Primary antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	192807312
NCBI GeneID	6597
NCBI Accession #	NP_001122316.1
NCBI GenBank Nucleotide #	NM_001128844.1
UniProt Accession #	P51532; Q3TKT4; Q8K1P7
Molecular Weight	1647
NCBI Official Full Name	transcription activator BRG1 isoform B
NCBI Official Synonym Full Names	SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily a, member 4
NCBI Official Symbol	SMARCA4
NCBI Official Synonym Symbols	BRG1; SNF2; SWI2; MRD16; RTPS2; BAF190; SNF2L4; SNF2LB; hSNF2b; BAF190A
NCBI Protein Information	transcription activator BRG1; SNF2-beta; SNF2-like 4; protein BRG-1; nuclear protein GRB1; brahma protein-like 1; BRM/SWI2-related gene 1; homeotic gene regulator; protein brahma homolog 1; BRG1-associated factor 190A; sucrose nonfermenting-like 4; ATP-de
UniProt Protein Name	Transcription activator BRG1
UniProt Gene Name	SMARCA4
UniProt Synonym Gene Names	BAF190A; BRG1; SNF2B; SNF2L4; BAF190A
UniProt Entry Name	SMCA4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/brg1-antibody-immunoprecipitation-immunofluorescence-immunohistochemistry-western-blot-elisa/223057>