

Rabbit Caspase 3 p17 Polyclonal Antibody | anti-CASP3 antibody

Rabbit anti-Caspase 3 p17 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA223653	Rabbit	Human, Mouse, Rat, Bovine, Monkey, Sheep	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Caspase 3 p17; Polyclonal Antibody; Rabbit anti-Caspase 3 p17 Antibody; CPP32; Caspase-3; CASP-3; Apopain; Cysteine protease CPP32; CPP-32; Protein Yama; SREBP cleavage activity 1; SCA-1; anti-CASP3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Monkey, Sheep
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-CASP3 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Caspase 3 p17. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Caspase 3 p17 staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of Caspase 3 p17 staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of Caspase 3 p17 expression in SP2/0 (A), Jurkat etoposide-treated (B), NIH3T3 staurosporine-treated (C), HeLa (D) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-CASP3 antibody

Primary antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	14790119
NCBI GeneID	836
NCBI Accession #	NP_004337.2
NCBI GenBank Nucleotide #	NM_004346.3
UniProt Accession #	P42574; P70677; P55213
Molecular Weight	31,608 Da
NCBI Official Full Name	caspase-3 preproprotein
NCBI Official Synonym Full Names	caspase 3, apoptosis-related cysteine peptidase
NCBI Official Symbol	CASP3
NCBI Official Synonym Symbols	CPP32; SCA-1; CPP32B
NCBI Protein Information	caspase-3; CASP-3; CPP-32; apopain; procaspase3; protein Yama; PARP cleavage protease; cysteine protease CPP32; SREBP cleavage activity 1; caspase 3, apoptosis-related cysteine protease
UniProt Protein Name	Caspase-3
UniProt Gene Name	CASP3
UniProt Synonym Gene Names	CPP32; CASP-3; CPP-32; SCA-1
UniProt Entry Name	CASP3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/caspase-3-p17-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/223653>