

Rabbit PGHS-2 Polyclonal Antibody | anti-PGHS-2 antibody

Rabbit anti-PGHS-2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA223733	Rabbit	Human, Mouse, Rat, Bovine, Pig, Sheep	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

PGHS-2; Polyclonal Antibody; Rabbit anti-PGHS-2 Antibody; COX2; Prostaglandin G/H synthase 2; Cyclooxygenase-2; COX-2; PHS II; Prostaglandin H2 synthase 2; PGH synthase 2; Prostaglandin-endoperoxide synthase 2; anti-PGHS-2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Pig, Sheep
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-PGHS-2 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human PGHS-2. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of PGHS-2 staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of PGHS-2 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of PGHS-2 expression in HeLa (A), Jurkat (B), Raw264.7 (C), rat lung (D) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-PGHS-2 antibody	Primary antibodies
--	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	4506265
NCBI GeneID	5743
NCBI Accession #	NP_000954.1
NCBI GenBank Nucleotide #	NM_000963.2
UniProt Accession #	P35354; Q05769; P35355
Molecular Weight	68,996 Da
NCBI Official Full Name	prostaglandin G/H synthase 2
NCBI Official Synonym Full Names	prostaglandin-endoperoxide synthase 2 (prostaglandin G/H synthase and cyclooxygenase)
NCBI Official Symbol	PTGS2
NCBI Official Synonym Symbols	COX2; COX-2; PHS-2; PGG/HS; PGHS-2; hCox-2; GRIPGHS
NCBI Protein Information	prostaglandin G/H synthase 2; PHS II; PGH synthase 2; cyclooxygenase 2b; prostaglandin H2 synthase 2
UniProt Protein Name	Prostaglandin G/H synthase 2
UniProt Gene Name	PTGS2
UniProt Synonym Gene Names	COX2; COX-2; PGH synthase 2; PGHS-2
UniProt Entry Name	PGH2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://stage.aaabiotech.com/polyclonal/pghs-2-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/223733>