

Rabbit SUMO1 Polyclonal Antibody | anti-SUMO1 antibody

Rabbit anti-SUMO1 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA223762	Rabbit	Human, Mouse, Rat, Bovine, Dog, Monkey, Pig	Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

SUMO1; Polyclonal Antibody; Rabbit anti-SUMO1 Antibody; SMT3C; SMT3H3; UBL1; Small ubiquitin-related modifier 1; SUMO-1; GAP-modifying protein 1; GMP1; SMT3 homolog 3; Sentrin; Ubiquitin-homology domain protein PIC1; Ubiquitin-like protein SMT3C; Smt3C; Ubiquitin-like protein UBL1; anti-SUMO1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Dog, Monkey, Pig
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-SUMO1 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human SUMO1. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of SUMO1 staining in A431 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of SUMO1 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of SUMO1 expression in MCF7 (A), A431 (B), HeLa (C), mouse kidney (D), rat kidney (E) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-SUMO1 antibody	Primary antibodies
---	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	54792065
NCBI GeneID	7341
NCBI Accession #	NP_001005781.1
NCBI GenBank Nucleotide #	NM_001005781.1
UniProt Accession #	P63165; P63166; Q5I0H3
Molecular Weight	11,557 Da
NCBI Official Full Name	small ubiquitin-related modifier 1 isoform a
NCBI Official Synonym Full Names	small ubiquitin-like modifier 1
NCBI Official Symbol	SUMO1
NCBI Official Synonym Symbols	DAP1; GMP1; PIC1; SMT3; UBL1; OFC10; SENP2; SMT3C; SMT3H3
NCBI Protein Information	small ubiquitin-related modifier 1; sentrin; SMT3 homolog 3; GAP modifying protein 1; ubiquitin-like protein UBL1; ubiquitin-like protein SMT3C; SMT3 suppressor of mif two 3 homolog 1; ubiquitin-homology domain protein PIC1
UniProt Protein Name	Small ubiquitin-related modifier 1
UniProt Gene Name	SUMO1
UniProt Synonym Gene Names	SMT3C; SMT3H3; UBL1; SUMO-1; GMP1; Smt3C
UniProt Entry Name	SUMO1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/sumo1-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/223762>