

Rabbit CKMT2 Polyclonal Antibody | anti-CKMT2 antibody

Rabbit anti-CKMT2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA223809	Rabbit	Human, Mouse, Rat, Bovine, Chicken, Dog, Pig	Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

CKMT2; Polyclonal Antibody; Rabbit anti-CKMT2 Antibody; Creatine kinase S-type; mitochondrial; Basic-type mitochondrial creatine kinase; Mib-CK; Sarcomeric mitochondrial creatine kinase; S-MtCK; anti-CKMT2 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Chicken, Dog, Pig
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-CKMT2 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CKMT2. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of CKMT2 staining in THP1 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of CKMT2 staining in human muscle formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of CKMT2 expression in THP1 (A), HeLa (B), Jurkat (C) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-CKMT2 antibody	Primary antibodies
---	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	153251984
NCBI GeneID	1160
NCBI Accession #	NP_001093205.1
NCBI GenBank Nucleotide #	NM_001099735.1
UniProt Accession #	P17540; Q6P8J7; P09605
Molecular Weight	47,504 Da
NCBI Official Full Name	creatine kinase S-type, mitochondrial
NCBI Official Synonym Full Names	creatine kinase, mitochondrial 2 (sarcomeric)
NCBI Official Symbol	CKMT2
NCBI Official Synonym Symbols	SMTCK
NCBI Protein Information	creatine kinase S-type, mitochondrial; S-MtCK; mib-CK; basic-type mitochondrial creatine kinase; sarcomeric mitochondrial creatine kinase
UniProt Protein Name	Creatine kinase S-type, mitochondrial
UniProt Gene Name	CKMT2
UniProt Synonym Gene Names	Mib-CK; S-MtCK
UniProt Entry Name	KCRS_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/ckmt2-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/223809>