Rabbit Alpha-synuclein Polyclonal Antibody | anti-SNCA antibody

Rabbit anti-Alpha-synuclein (pS129) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA223913	**Rabbit**	**Human, Mouse, Rat, Bovine, Monkey, Sheep**	**Immunofluorescence, Immunohistochemistry, Western Blot**

PRODUCT SYNONYMS

Alpha-synuclein; Polyclonal Antibody; Rabbit anti-Alpha-synuclein (pS129) Antibody; NACP; PARK1; Non-A beta component of AD amyloid; Non-A4 component of amyloid precursor; anti-SNCA antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Monkey, Sheep
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-SNCA antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Alpha-synuclein (pS129). The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.


IMAGES



IF (Immunofluorescence)
Immunofluorescent analysis of Alpha-synuclein (pS129) staining in HEK293T cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohiostchemistry)
Immunohistochemical analysis of Alpha-synuclein (pS129) staining in human prostate cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)
Western blot analysis of Alpha-synuclein (pS129) expression in HEK293T (A), mouse brain (B) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-SNCA antibody	Primary antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	4507109
NCBI GeneID	6622
NCBI Accession #	NP_000336.1
NCBI GenBank Nucleotide #	NM_000345.3
UniProt Accession #	P37840; O55042; P37377
Molecular Weight	14,460 Da
NCBI Official Full Name	alpha-synuclein isoform NACP140
NCBI Official Synonym Full Names	synuclein, alpha (non A4 component of amyloid precursor)
NCBI Official Symbol	SNCA
NCBI Official Synonym Symbols	PD1; NACP; PARK1; PARK4
NCBI Protein Information	alpha-synuclein; synuclein alpha-140; non A-beta component of AD amyloid
UniProt Protein Name	Alpha-synuclein
UniProt Gene Name	**SNCA**
UniProt Synonym Gene Names	NACP; PARK1; NACP
UniProt Entry Name	SYUA_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: https://stage.aaabiotech.com/polyclonal/alpha-synuclein-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/223913