

Rabbit TBP Polyclonal Antibody | anti-TBP antibody

Rabbit anti-TBP Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA223916	Rabbit	Human, Mouse, Rat, Bovine, Chicken, Pig, Rabbit, Zebrafish	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

TBP; Polyclonal Antibody; Rabbit anti-TBP Antibody; GTF2D1; TF2D; TFIID; TATA-box-binding protein; TATA sequence-binding protein; TATA-binding factor; TATA-box factor; Transcription initiation factor TFIID TBP subunit; anti-TBP antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Chicken, Pig, Rabbit, Zebrafish
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-TBP antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human TBP. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of TBP staining in HEK293T cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of TBP staining in human prostate cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of TBP expression in HEK293T (A), NIH3T3 (B) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-TBP antibody

[Primary antibodies](#)

NCBI AND UNIPROT INFORMATION

NCBI GI #	285026520
NCBI GeneID	6908
NCBI Accession #	NP_001165556.1
NCBI GenBank Nucleotide #	NM_001172085.1
UniProt Accession #	P20226 ; P29037
Molecular Weight	37,698 Da
NCBI Official Full Name	TATA-box-binding protein isoform 2
NCBI Official Synonym Full Names	TATA box binding protein
NCBI Official Symbol	TBP
NCBI Official Synonym Symbols	HDL4; GTF2D; SCA17; TFIID; GTF2D1
NCBI Protein Information	TATA-box-binding protein; TATA-box factor; TATA sequence-binding protein; TATA-box binding protein N-terminal domain; transcription initiation factor TFIID TBP subunit
UniProt Protein Name	TATA-box-binding protein
UniProt Gene Name	TBP
UniProt Synonym Gene Names	GTF2D1; TF2D; TFIID
UniProt Entry Name	TBP_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/tbp-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/223916>