

Rabbit SF1 Polyclonal Antibody | anti-SF1 antibody

Rabbit anti-SF1 (pS82) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA223938	Rabbit	Human, Mouse, Rat, Bovine, Rabbit, Zebrafish	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

SF1; Polyclonal Antibody; Rabbit anti-SF1 (pS82) Antibody; ZFM1; ZNF162; Splicing factor 1; Mammalian branch point-binding protein; BBP; mBBP; Transcription factor ZFM1; Zinc finger gene in MEN1 locus; Zinc finger protein 162; anti-SF1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Rabbit, Zebrafish
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-SF1 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human SF1 (pS82). The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of SF1 (pS82) staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of SF1 (pS82) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of SF1 (pS82) expression in SHSY5Y (A), Jurkat (B) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-SF1 antibody

[Primary antibodies](#)

NCBI AND UNIPROT INFORMATION

NCBI GI #	295842330
NCBI GeneID	7536
NCBI Accession #	NP_001171501.1
NCBI GenBank Nucleotide #	NM_001178030.1
UniProt Accession #	Q15637 ; Q64213
Molecular Weight	68,330 Da
NCBI Official Full Name	splicing factor 1 isoform 6
NCBI Official Synonym Full Names	splicing factor 1
NCBI Official Symbol	SF1
NCBI Official Synonym Symbols	BBP; MBBP; ZFM1; ZNF162; D11S636; ZCCHC25
NCBI Protein Information	splicing factor 1; zinc finger protein 162; transcription factor ZFM1; zinc finger gene in MEN1 locus; mammalian branch point-binding protein
UniProt Protein Name	Splicing factor 1
UniProt Gene Name	SF1
UniProt Synonym Gene Names	ZFM1; ZNF162; BBP; mBBP
UniProt Entry Name	SF01_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/sf1-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/223938>