

Rabbit MKK4 Polyclonal Antibody | anti-MKK4 antibody

Rabbit anti-MKK4 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA223959	Rabbit	Human, Mouse, Rat, Bovine, Zebrafish	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

MKK4; Polyclonal Antibody; Rabbit anti-MKK4 Antibody; JNKK1; MEK4; PRKMK4; SEK1; SERK1; SKK1; Dual specificity mitogen-activated protein kinase kinase 4; MAP kinase kinase 4; MAPKK 4; JNK-activating kinase 1; MAPK/ERK kinase 4; MEK 4; SAPK/ERK kinase 1; Stress-activated protein kinase kinase 1; SAPK kinase 1; SAPKK-1; SAPKK1; c-Jun N-terminal kinase kinase 1; JNKK; anti-MKK4 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Zebrafish
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-MKK4 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human MKK4. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of MKK4 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of MKK4 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of MKK4 expression in HepG2 (A), K562 (B), HeLa (C) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-MKK4 antibody	Primary antibodies
--	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	4506889
NCBI GeneID	6416
NCBI Accession #	NP_003001.1
NCBI GenBank Nucleotide #	NM_003010.2
UniProt Accession #	P45985; P47809
Molecular Weight	44,288 Da
NCBI Official Full Name	dual specificity mitogen-activated protein kinase kinase 4
NCBI Official Synonym Full Names	mitogen-activated protein kinase kinase 4
NCBI Official Symbol	MAP2K4
NCBI Official Synonym Symbols	JNKK; MEK4; MKK4; SEK1; JNKK1; SERK1; MAPKK4; PRKMK4; SAPKK1; SAPKK-1
NCBI Protein Information	dual specificity mitogen-activated protein kinase kinase 4; MEK 4; MAPKK 4; SAPK kinase 1; MAPK/ERK kinase 4; SAPK/ERK kinase 1; MAP kinase kinase 4; JNK-activated kinase 1; JNK activating kinase 1; JNK-activating kinase 1; c-Jun N-terminal kinase kinase
UniProt Protein Name	Dual specificity mitogen-activated protein kinase kinase 4
UniProt Gene Name	MAP2K4
UniProt Synonym Gene Names	JNKK1; MEK4; MKK4; PRKMK4; SEK1; SERK1; SKK1; MAP kinase kinase 4; MAPKK 4; MEK 4; SEK1; SAPK kinase 1; SAPKK-1; SAPKK1; JNKK
UniProt Entry Name	MP2K4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.