

Rabbit ZNF265 Polyclonal Antibody | anti-ZNF265 antibody

Rabbit anti-ZNF265 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA223977	Rabbit	Human, Mouse, Rat, Chicken, Dog, Monkey, Pig, Rabbit	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

ZNF265; Polyclonal Antibody; Rabbit anti-ZNF265 Antibody; ZIS; Zinc finger Ran-binding domain-containing protein 2; Zinc finger protein 265; Zinc finger; splicing; anti-ZNF265 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Chicken, Dog, Monkey, Pig, Rabbit
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-ZNF265 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human ZNF265. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of ZNF265 staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of ZNF265 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of ZNF265 expression in HepG2 (A), MCF7 (B) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-ZNF265 antibody

[Primary antibodies](#)

NCBI AND UNIPROT INFORMATION

NCBI GI #	42741682
NCBI GeneID	9406
NCBI Accession #	NP_005446.2
NCBI GenBank Nucleotide #	NM_005455.4
UniProt Accession #	O95218; Q9R020; O35986
Molecular Weight	37,404 Da
NCBI Official Full Name	zinc finger Ran-binding domain-containing protein 2 isoform 2
NCBI Official Synonym Full Names	zinc finger, RAN-binding domain containing 2
NCBI Official Symbol	ZRANB2
NCBI Official Synonym Symbols	ZIS; ZIS1; ZIS2; ZNF265
NCBI Protein Information	zinc finger Ran-binding domain-containing protein 2; zinc finger, splicing; zinc-finger, splicing; zinc finger protein 265; zinc-finger protein 265
UniProt Protein Name	Zinc finger Ran-binding domain-containing protein 2
UniProt Gene Name	ZRANB2
UniProt Synonym Gene Names	ZIS; ZNF265
UniProt Entry Name	ZRAB2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/znf265-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/223977>