

Rabbit S6K1 Polyclonal Antibody | anti-S6K1 antibody

Rabbit anti-S6K1 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA223978	Rabbit	Human, Mouse, Rat, Bovine, Rabbit	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

S6K1; Polyclonal Antibody; Rabbit anti-S6K1 Antibody; STK14A; Ribosomal protein S6 kinase beta-1; S6K-beta-1; 70 kDa ribosomal protein S6 kinase 1; P70S6K1; p70-S6K 1; Ribosomal protein S6 kinase I; Serine/threonine-protein kinase 14A; p70 ribosomal S6 kinase alpha; p70 S6 kinase alpha; p70 S6K-alpha; p70 S6KA; anti-S6K1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Rabbit
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-S6K1 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human S6K1. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of S6K1 staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of S6K1 staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)
Western blot analysis of S6K1 expression in HepG2 (A), Jurkat (B) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-S6K1 antibody	Primary antibodies
--	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	440546416
NCBI GeneID	6198
NCBI Accession #	NP_001258989.1
NCBI GenBank Nucleotide #	NM_001272060.1
UniProt Accession #	P23443; Q8BSK8; P67999
Molecular Weight	59,140 Da
NCBI Official Full Name	ribosomal protein S6 kinase beta-1 isoform e
NCBI Official Synonym Full Names	ribosomal protein S6 kinase, 70kDa, polypeptide 1
NCBI Official Symbol	RPS6KB1
NCBI Official Synonym Symbols	S6K; PS6K; S6K1; STK14A; p70-S6K; p70 S6KA; p70-alpha; S6K-beta-1; p70(S6K)-alpha
NCBI Protein Information	ribosomal protein S6 kinase beta-1; p70 S6 kinase, alpha; ribosomal protein S6 kinase I; serine/threonine kinase 14 alpha; serine/threonine-protein kinase 14A
UniProt Protein Name	Ribosomal protein S6 kinase beta-1
UniProt Gene Name	RPS6KB1
UniProt Synonym Gene Names	STK14A; S6K-beta-1; S6K1; P70S6K1; p70-S6K 1; p70 S6 kinase alpha; p70 S6K-alpha; p70 S6KA
UniProt Entry Name	KS6B1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.