

Rabbit MKK6 Polyclonal Antibody | anti-MKK6 antibody

Rabbit anti-MKK6 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA223987	Rabbit	Human, Mouse, Rat, Bovine, Monkey	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

MKK6; Polyclonal Antibody; Rabbit anti-MKK6 Antibody; MEK6; PRKMK6; SKK3; Dual specificity mitogen-activated protein kinase kinase 6; MAP kinase kinase 6; MAPKK 6; MAPK/ERK kinase 6; MEK 6; Stress-activated protein kinase kinase 3; SAPK kinase 3; SAPKK-3; SAPKK3; anti-MKK6 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Monkey
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-MKK6 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human MKK6. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of MKK6 staining in Hela cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of MKK6 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)
Western blot analysis of MKK6 expression in Hela (A), HEK293T (B) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-MKK6 antibody	Primary antibodies
--	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	14589900
NCBI GeneID	5608
NCBI Accession #	NP_002749.2
NCBI GenBank Nucleotide #	NM_002758.3
UniProt Accession #	P52564; P70236
Molecular Weight	37,492 Da
NCBI Official Full Name	dual specificity mitogen-activated protein kinase kinase 6
NCBI Official Synonym Full Names	mitogen-activated protein kinase kinase 6
NCBI Official Symbol	MAP2K6
NCBI Official Synonym Symbols	MEK6; MKK6; MAPKK6; PRKMK6; SAPKK3; SAPKK-3
NCBI Protein Information	dual specificity mitogen-activated protein kinase kinase 6; MEK 6; MAPKK 6; SAPK kinase 3; MAPK/ERK kinase 6; stress-activated protein kinase kinase 3; protein kinase, mitogen-activated, kinase 6 (MAP kinase kinase 6)
UniProt Protein Name	Dual specificity mitogen-activated protein kinase kinase 6
UniProt Gene Name	MAP2K6
UniProt Synonym Gene Names	MEK6; MKK6; PRKMK6; SKK3; MAP kinase kinase 6; MAPKK 6; MEK 6; SAPK kinase 3; SAPKK-3; SAPKK3
UniProt Entry Name	MP2K6_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.