

Rabbit CD29 Polyclonal Antibody | anti-CD29 antibody

Rabbit anti-CD29 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA224016	Rabbit	Human, Mouse, Rat, Bovine, Chicken, Pig, Sheep	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

CD29; Polyclonal Antibody; Rabbit anti-CD29 Antibody; FNRB; MDF2; MSK12; Integrin beta-1; Fibronectin receptor subunit beta; Glycoprotein IIa; GPIIa; VLA-4 subunit beta; anti-CD29 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Chicken, Pig, Sheep
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-CD29 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human CD29. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of CD29 staining in NIH3T3 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of CD29 staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)
 Western blot analysis of CD29 expression in HepG2 (A), Jurkat (B) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-CD29 antibody	Primary antibodies
--	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	19743813
NCBI GeneID	3688
NCBI Accession #	NP_002202.2
NCBI GenBank Nucleotide #	NM_002211.3
UniProt Accession #	P05556; P09055; P49134
Molecular Weight	88,415 Da
NCBI Official Full Name	integrin beta-1 isoform 1A
NCBI Official Synonym Full Names	integrin, beta 1 (fibronectin receptor, beta polypeptide, antigen CD29 includes MDF2, MSK12)
NCBI Official Symbol	ITGB1
NCBI Official Synonym Symbols	CD29; FNRB; MDF2; VLAB; GPIIA; MSK12; VLA-BETA
NCBI Protein Information	integrin beta-1; glycoprotein IIa; integrin VLA-4 beta subunit; very late activation protein, beta polypeptide
UniProt Protein Name	Integrin beta-1
UniProt Gene Name	ITGB1
UniProt Synonym Gene Names	FNRB; MDF2; MSK12; GPIIA
UniProt Entry Name	ITB1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.