

Rabbit Calnexin Polyclonal Antibody | anti-CDK1 antibody

Rabbit anti-Calnexin Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA224078	Rabbit	Human, Mouse, Rat, Bovine, Dog	Immunoprecipitation, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Calnexin; Polyclonal Antibody; Rabbit anti-Calnexin Antibody; IP90; Major histocompatibility complex class I antigen-binding protein p88; p90; anti-CDK1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Dog
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-CDK1 antibody	IP (Immunoprecipitation), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Calnexin. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of CDC2 staining in K562 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of CDC2 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of CDC2 expression in HeLa (A), K562 (B), MCF7 (C), A431 (D) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-CDK1 antibody	Primary antibodies
--	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	4502709
NCBI GeneID	983
NCBI Accession #	NP_001777.1
NCBI GenBank Nucleotide #	NM_001786.4
UniProt Accession #	P06493; P11440; P39951
Molecular Weight	34,095 Da
NCBI Official Full Name	cyclin-dependent kinase 1 isoform 1
NCBI Official Synonym Full Names	cyclin-dependent kinase 1
NCBI Official Symbol	CDK1
NCBI Official Synonym Symbols	CDC2; CDC28A; P34CDC2
NCBI Protein Information	cyclin-dependent kinase 1; p34 protein kinase; cell cycle controller CDC2; cell division protein kinase 1; cell division control protein 2 homolog; cell division cycle 2, G1 to S and G2 to M
UniProt Protein Name	Cyclin-dependent kinase 1
UniProt Gene Name	CDK1
UniProt Synonym Gene Names	CDC2; CDC28A; CDKN1; P34CDC2; CDK1
UniProt Entry Name	CDK1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/calnexin-antibody-immunoprecipitation-immunofluorescence-immunohistochemistry-western-blot-elisa/224078>