

Rabbit Lamin A/C Polyclonal Antibody | anti-SMAD4 antibody

Rabbit anti-Lamin A/C Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA224094	Rabbit	Human, Mouse, Rat, Bovine, Monkey, Pig	Immunoprecipitation, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

Lamin A/C; Polyclonal Antibody; Rabbit anti-Lamin A/C Antibody; LMN1; Prelamin-A/C; anti-SMAD4 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Monkey, Pig
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-SMAD4 antibody	IP (Immunoprecipitation), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Lamin A/C. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of SMAD4 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of SMAD4 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of SMAD4 expression in HepG2 (A), HeLa (B), NIH3T3 (C), COS7 (D) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-SMAD4 antibody	Primary antibodies
---	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	4885457
NCBI GeneID	4089
NCBI Accession #	NP_005350.1
NCBI GenBank Nucleotide #	NM_005359.5
UniProt Accession #	Q13485; P97471; O70437
Molecular Weight	60,439 Da
NCBI Official Full Name	mothers against decapentaplegic homolog 4
NCBI Official Synonym Full Names	SMAD family member 4
NCBI Official Symbol	SMAD4
NCBI Official Synonym Symbols	JIP; DPC4; MADH4; MYHRS
NCBI Protein Information	mothers against decapentaplegic homolog 4; MAD homolog 4; SMAD, mothers against DPP homolog 4; deleted in pancreatic carcinoma locus 4; deletion target in pancreatic carcinoma 4; mothers against decapentaplegic, Drosophila, homolog of, 4
UniProt Protein Name	Mothers against decapentaplegic homolog 4
UniProt Gene Name	SMAD4
UniProt Synonym Gene Names	DPC4; MADH4; MAD homolog 4; Mothers against DPP homolog 4; SMAD 4; Smad4; hSMAD4
UniProt Entry Name	SMAD4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/lamin-a-c-antibody-immunoprecipitation-immunofluorescence-immunohistochemistry-western-blot-elisa/224094>