

Rabbit MKK4 Polyclonal Antibody | anti-MKK4 antibody

Rabbit anti-MKK4 (pS80) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA224103	Rabbit	Human, Mouse, Rat, Bovine, Zebrafish	Immunoprecipitation, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

MKK4; Polyclonal Antibody; Rabbit anti-MKK4 (pS80) Antibody; JNKK1; MEK4; PRKMK4; SEK1; SERK1; SKK1; Dual specificity mitogen-activated protein kinase kinase 4; MAP kinase kinase 4; MAPKK 4; JNK-activating kinase 1; MAPK/ERK kinase 4; MEK 4; SAPK/ERK kinase 1; Stress-activated protein kinase kinase 1; SAPK kinase 1; SAPKK-1; SAPKK1; c-Jun N-terminal kinase kinase 1; JNKK; anti-MKK4 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Zebrafish
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-MKK4 antibody	IP (Immunoprecipitation), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human MKK4. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of STAT1 (pS727) staining in NIH3T3 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of STAT1 (pS727) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of STAT1 (pS727) expression in A431 (A), NIH3T3 insulin-treated (B), Jurkat pervanadate-treated (C) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-MKK4 antibody

Primary antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	6274552
NCBI GeneID	6772
NCBI Accession #	NP_009330.1
NCBI GenBank Nucleotide #	NM_007315.3
UniProt Accession #	P42224; P42225
Molecular Weight	87,335 Da
NCBI Official Full Name	signal transducer and activator of transcription 1-alpha/beta isoform alpha
NCBI Official Synonym Full Names	signal transducer and activator of transcription 1, 91kDa
NCBI Official Symbol	STAT1
NCBI Official Synonym Symbols	CANDF7; ISGF-3; STAT91
NCBI Protein Information	signal transducer and activator of transcription 1-alpha/beta; transcription factor ISGF-3 components p91/p84; signal transducer and activator of transcription-1
UniProt Protein Name	Signal transducer and activator of transcription 1-alpha/beta
UniProt Gene Name	STAT1
UniProt Entry Name	STAT1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/mkk4-antibody-immunoprecipitation-immunofluorescence-immunohistochemistry-western-blot-elisa/224103>