

Rabbit Flotillin 2 Polyclonal Antibody | anti-FLOT2 antibody

Flotillin 2 Antibody

CATALOG # AAA225078	HOST Rabbit	APPLICATIONS Western Blot, Immunohistochemistry, Immunocytochemistry, Immunofluorescence
-------------------------------	-----------------------	--

PRODUCT SYNONYMS

Flotillin 2; Polyclonal Antibody; Flotillin 2 Antibody; Affinity Purified Rabbit Polyclonal Flotillin 2 Antibody; Polyclonal Flotillin 2 antibody; Anti-Flotillin 2 antibody; M17S1 antibody; epidermal surface antigen antibody; Flotillin 2 (epidermal surface antigen 1) antibody; Flotillin -2; ESA1 antibody; ECS1 antibody; ECS-1 antibody; Flotillin 2 antibody; FLOT2 antibody; flotillin-2 antibody; ESA antibody; Flotillin -2 antibody; membrane component chromosome 17 surface marker 1 (35kD protein identified by antibody ECS-1) antibody; flotillin 2 antibody; membrane component chromosome 17 surface marker 1 antibody; anti-FLOT2 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen-Affinity Chromatography
Form/Format	Supplied as a concentrated solution containing 1X PBS, 20% Glycerol (pH 7.0). 0.01% Thimerosal was added as a preservative.
Concentration	1 mg/ml (varies by lot)
Sequence Length	448
Applicable Applications for anti-FLOT2 antibody	WB (Western Blot), IHC (Immunohistochemistry), ICC (Immunocytochemistry), IF (Immunofluorescence)
Immunogen	Flotillin 2 antibody was raised in rabbit using a synthetic peptide corresponding to a region within amino acids 365 and 428 of Flotillin-2
Cross-Reactivity	Human
Preparation and Storage	Store at 4 degree C for short term storage. Aliquot and store at -20 degree C for long term storage. Avoid repeated freeze/thaw cycles.

IMAGES




WB (Western Blot)

Western blot analysis of 30 ug of whole cell lysate (A:HeLa S3) using a 10% SDS PAGE gel and Flotillin 2 antibody at a dilution of 1:1000



IHC (Immunohistochemistry)

Immunohistochemical staining of paraffin-embedded Lung CA using Flotillin 2 antibody at a dilution of 1:100



IF (Immunofluorescence)

Immunofluorescence analysis of paraformaldehyde-fixed HeLaS3, using Flotillin-2 antibody at 1:100 dilution.

ADDITIONAL INFORMATION

Related Product Information for anti-FLOT2 antibody	Caveolae are small domains on the inner cell membrane involved in vesicular trafficking and signal transduction. This gene encodes a caveolae-associated, integral membrane protein, which is thought to function in neuronal signaling.
Product Categories/Family for anti-FLOT2 antibody	Signal Transduction

NCBI AND UNIPROT INFORMATION

NCBI GI #	442616360
NCBI GeneID	32425
NCBI GenBank Nucleotide #	NP_001259553.1
Molecular Weight	47 kDa
NCBI Official Full Name	flotillin 2, isoform J
NCBI Official Synonym Full Names	Flotillin 2
NCBI Official Symbol	Flo2
NCBI Official Synonym Symbols	CG11547; CG14409; CG32593; DmelCG32593; DmFlo2; Flo; Flo-2; flo2; FLO2; FLO[Dm]-2; mdcds_54791; reggie-1
NCBI Protein Information	flotillin 2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/flotillin-2-antibody-western-blot-immunohistochemistry-immunocytochemistry-immunofluorescence-elisa/225078>