

Rabbit ULK2 Polyclonal Antibody | anti-ULK2 antibody

ULK2 Antibody

CATALOG # AAA225539	HOST Rabbit	APPLICATIONS Western Blot, Immunohistochemistry, Immunocytochemistry, Immunofluorescence
-------------------------------	-----------------------	--


PRODUCT SYNONYMS

ULK2; Polyclonal Antibody; ULK2 Antibody; Affinity Purified Rabbit Polyclonal ULK2 Antibody; Polyclonal ULK2 antibody; Anti-ULK2 antibody; ULK 2; ULK 2 antibody; ULK-2; unc-51-like kinase 2 (C. elegans) antibody; KIAA0623 antibody; ULK2 antibody; ATG1B antibody; serine/threonine-protein kinase ULK2 antibody; Unc51.2 antibody; ULK-2 antibody; anti-ULK2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen-Affinity Chromatography
Form/Format	Supplied as a concentrated solution containing 0.1M Tris, 0.1M Glycine, 10% Glycerol (pH 7.0). 0.01% Thimerosal was added as a preservative.
Concentration	1 mg/ml (varies by lot)
Sequence Length	133
Applicable Applications for anti-ULK2 antibody	WB (Western Blot), IHC (Immunohistochemistry), ICC (Immunocytochemistry), IF (Immunofluorescence)
Cross-Reactivity	Human
Preparation and Storage	Store at 4 degree C for short term storage. Aliquot and store at -20 degree C for long term storage. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescence analysis of paraformaldehyde-fixed HeLa, using ULK2 antibody at 1:200 dilution.




WB (Western Blot)

Western blot analysis of 30 ug of whole cell lysate (A: NIH-3T3) using a 7.5% SDS PAGE gel and ULK2 antibody at a dilution of 1:1000



IHC (Immunohistochemistry)

Immunohistochemical staining of paraffin-embedded MCF7 xenograft using ULK2 antibody at a dilution of 1:500



WB (Western Blot)

Western blot analysis of 30 ug of whole cell lysate (A: NT2D1; B: PC-3) using a 7.5% SDS PAGE gel and ULK2 antibody at a dilution of 1:1000

ADDITIONAL INFORMATION

Related Product Information for anti-ULK2 antibody	This gene encodes a protein that is similar to a serine/threonine kinase in C. elegans which is involved in axonal elongation. The structure of this protein is similar to the C. elegans protein in that both proteins have an N-terminal kinase domain, a central proline/serine rich domain, and a C-terminal domain. The gene is located within the Smith-Magenis syndrome region on chromosome 17.
Product Categories/Family for anti-ULK2 antibody	Proteases; Inhibitors; & Enzymes

NCBI AND UNIPROT INFORMATION

NCBI GI #	958277893
Molecular Weight	113 kDa
NCBI Official Full Name	ULK2, partial

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/ulk2-antibody-western-blot-immunohistochemistry-immunocytochemistry-immunofluorescence-elisa/225539>