

Rabbit TRIM32 Polyclonal Antibody | anti-TRIM32 antibody

TRIM32 Antibody

CATALOG # AAA225610	HOST Rabbit	APPLICATIONS Western Blot, Immunohistochemistry, Immunocytochemistry, Immunofluorescence
-------------------------------	-----------------------	--

PRODUCT SYNONYMS

TRIM32; Polyclonal Antibody; TRIM32 Antibody; Affinity Purified Rabbit Polyclonal TRIM32 Antibody; Polyclonal TRIM32 antibody; Anti-TRIM32 antibody; TATIP antibody; HT2A antibody; tripartite motif-containing protein 32 antibody; 72 kDa Tat-interacting protein antibody; TRIM 32; TRIM32 antibody; TRIM-32 antibody; Zinc-finger protein HT2A antibody; TRIM 32 antibody; Zinc finger protein HT2A antibody; LGMD2H antibody; TRIM-32; BBS11 antibody; tripartite motif-containing 32 antibody; anti-TRIM32 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen-Affinity Chromatography
Form/Format	Supplied as a concentrated solution containing 0.1M Tris, 0.1M Glycine, 20% Glycerol (pH 7.0). 0.01% Thimerosal was added as a preservative.
Concentration	0.74 mg/ml (varies by lot)
Sequence Length	653
Applicable Applications for anti-TRIM32 antibody	WB (Western Blot), IHC (Immunohistochemistry), ICC (Immunocytochemistry), IF (Immunofluorescence)
Cross-Reactivity	Human
Preparation and Storage	Store at 4 degree C for short term storage. Aliquot and store at -20 degree C for long term storage. Avoid repeated freeze/thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemical staining of paraffin-embedded H441 Xenograft using TRIM32 antibody at a dilution of 1:100



WB (Western Blot)

Western blot analysis of 30 ug of whole cell lysate (A: 293T) using a 7.5% SDS PAGE gel and TRIM32 antibody at a dilution of 1:1000



IF (Immunofluorescence)

Immunofluorescence analysis of paraformaldehyde-fixed A431, using TRIM32 antibody at 1:500 dilution.

ADDITIONAL INFORMATION

Related Product Information for anti-TRIM32 antibody	The protein encoded by this gene is a member of the tripartite motif family. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. The protein localizes to cytoplasmic bodies. The protein has also been localized to the nucleus, where it interacts with the activation domain of the HIV-1 Tat protein. The Tat protein activates transcription of HIV-1 genes.
Product Categories/Family for anti-TRIM32 antibody	DNA & RNA

NCBI AND UNIPROT INFORMATION

NCBI GI #	823670132
Molecular Weight	72 kDa
NCBI Official Full Name	TRIM32, partial

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/trim32-antibody-western-blot-immunohistochemistry-immunocytochemistry-immunofluorescence-elisa/225610>