

Rabbit anti-Human, Dog EXOSC6 Polyclonal Antibody | anti-EXOSC6 antibody

EXOSC6 Antibody

CATALOG # AAA225820	HOST Rabbit	REACTIVITY Human, Dog	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	---------------------------------	---

PRODUCT SYNONYMS

EXOSC6; Polyclonal Antibody; EXOSC6 Antibody; Rabbit Polyclonal EXOSC6 Antibody raised against the N terminal of EXOSC6; Polyclonal EXOSC6 antibody; Anti-EXOSC6 antibody; EXOSC 6 antibody; EXOSC 6; EXOSC-6 antibody; Exosome Component 6 antibody; EXOSC-6; anti-EXOSC6 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Dog
Clonality	Polyclonal
Specificity	EXOSC6 antibody was raised against the N terminal of EXOSC6
Purity/Purification	Total IgG Protein A purified
Form/Format	Liquid; in 1x PBS buffer with 0.09% (w/v) Sodium Azide and 2% Sucrose.
Concentration	1 mg/ml (varies by lot)
Applicable Applications for anti-EXOSC6 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	EXOSC6 antibody was raised using the N terminal of EXOSC6 corresponding to a region with amino acids MPG D H R R I R G P E E S Q P P Q L Y A A D E E E A P G T R D P T R L R P V Y A R A G L L S Q A K
Cross-Reactivity	Human, Dog
Preparation and Storage	Store at 2-8 degree C for short periods. For longer periods of storage, store at -20 degree C. Avoid repeat freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

EXOSC6 antibody was used for immunohistochemistry at a concentration of 4-8 ug/ml to stain Hepatocytes (arrows) in Human Liver. Magnification is at 400X



WB (Western Blot)

EXOSC6 antibody used at 5 ug/ml to detect target protein.



IHC (Immunohistochemistry)

EXOSC6 antibody was used for immunohistochemistry at a concentration of 4-8 ug/ml to stain Skeletal muscle cells (arrows) in Human Muscle. Magnification is at 400X.

ADDITIONAL INFORMATION

Related Product Information for anti-EXOSC6 antibody	EXOSC6 constitutes one of the subunits of the multisubunit particle called exosome, which mediates mRNA degradation. The composition of human exosome is similar to its yeast counterpart. This protein is homologous to the yeast Mtr3 protein. Its exact function is not known, however, it has been shown using a cell-free RNA decay system that the exosome is required for rapid degradation of unstable mRNAs containing AU-rich elements (AREs), but not for poly(A) shortening.
Product Categories/Family for anti-EXOSC6 antibody	DNA & RNA

NCBI AND UNIPROT INFORMATION

Molecular Weight	30 kDa (MW of target protein)
------------------	-------------------------------

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/exosc6-antibody-immunohistochemistry-western-blot-elisa/225820>