

Rhod-FF AM Biochemical

Rhod-FF AM

CATALOG #
AAA227282


PRODUCT SYNONYMS

Rhod-FF AM; Rhod-FF AM biochemical

PRODUCT SPECIFICATIONS


Purity/Purification	98%
Solubility	DMSO: Soluble <1mg/ml refers to the product slightly soluble or insoluble
Preparation and Storage	Powder: -20 degree C for 3 years In solvent: -80 degree C for 2 years

IMAGES



Structure

Modified MMAF-C5-COOH, CAS 1404071-65-3



Structure

Modified MMAF-C5-COOH, CAS 1404071-65-3

ADDITIONAL INFORMATION

Related Product Information for Rhod-FF AM biochemical	Rhod-FF AM is a cell-permeant acetoxymethyl ester of the fluorescent calcium indicator rhod-FF. As rhod-FF AM enters cells, it is hydrolyzed by intracellular esterases to produce rhod-FF. Rhod-FF is a difluorinated analog of the cell-impermeant calcium indicator rhod-2. It has a very low affinity for calcium ($K_d = 320 \mu M$). Low affinity calcium indicators are particularly useful for studying compartments with high concentrations of calcium, such as endoplasmic reticulum, where high affinity dyes will be insensitive to luminal fluctuations. Rhod-FF has excitation/emission maxima of approximately 552 and 580 nm, respectively.
References	- Polson AG, et al. Antibody-drug conjugates for the treatment of non-Hodgkin's lymphoma: target and linker-drug selection [published correction appears in Cancer Res. 2010 Feb 1;70(3):1275. Slaga, Dion S [added]]. Cancer Res. 2009;69(6):2358-2364. - Doronina SO, et al. Enhanced activity of monomethylauristatin F through monoclonal antibody delivery: effects of linker technology on efficacy and toxicity. Bioconjug Chem. 2006;17(1):114-124.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

