

H-D-Phe-Pip-Arg-pNA acetate Biochemical | H-D-Phe-Pip-Arg-pNA biochemical

H-D-Phe-Pip-Arg-pNA acetate

CATALOG #
AAA227485


PRODUCT SYNONYMS

H-D-Phe-Pip-Arg-pNA acetate; H-D-Phe-Pip-Arg-pNA biochemical


PRODUCT SPECIFICATIONS

Purity/Purification	98%
CAS	115388-96-0
Solubility	<1mg/ml refers to the product slightly soluble or insoluble
Preparation and Storage	Powder: -20 degree C for 3 years In solvent: -80 degree C for 2 years

IMAGES



Structure
A 410099.1, amine-Boc, CAS 1613552-03-6



Structure
A 410099.1, amine-Boc, CAS 1613552-03-6

ADDITIONAL INFORMATION

Related Product Information for H-D-Phe-Pip-Arg-pNA biochemical	H-D-Phe-Pip-Arg-pNA (S-2238) acetate, a chromogenic substrate, is patterned after the N-terminal portion of the A alpha chain of fibrinogen, which is the natural substrate of thrombin. H-D-Phe-Pip-Arg-pNA acetate is specific for thrombin and is used to measure antithrombin-heparin cofactor (AT-III). The AT-III assay using H-D-Phe-Pip-Arg-pNA acetate is sensitive, accurate, and easy to perform.
References	1. Christopher P Tinworth, et al. PROTAC-Mediated Degradation of Bruton's Tyrosine Kinase Is Inhibited by Covalent Binding. ACS Chem Biol.2019 Mar 15;14(3):342-347

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/biochemical/h-d-phe-pip-arg-pna-acetate/227485>