

Boc-NH-PPG2 biochemical

Boc-NH-PPG2

CATALOG #
AAA227585


PRODUCT SYNONYMS

Boc-NH-PPG2; Boc-NH-PPG2 biochemical


PRODUCT SPECIFICATIONS

Purity/Purification	98%
CAS	1312905-31-9
Solubility	<1mg/ml refers to the product slightly soluble or insoluble
In vitro	PROTACs contain two different ligands connected by a linker; one is a ligand for an E3 ubiquitin ligase and the other is for the target protein. PROTACs exploit the intracellular ubiquitin-proteasome system to selectively degrade target proteins[1].
Preparation and Storage	Powder: -20 degree C for 3 years In solvent: -80 degree C for 2 years

IMAGES



Structure
Ficusonolide, CAS 1800503-81-4



Structure
Ficusonolide, CAS 1800503-81-4

ADDITIONAL INFORMATION

Related Product Information for Boc-NH-PPG2 biochemical	Boc-NH-PPG2 is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.
References	1. Din AU, et al. Antidiabetic Activity of Ficusonolide, a Triterpene Lactone from Ficus foveolata (Wall. ex Miq.): In Vitro, In Vivo, and In Silico Approaches. ACS Omega. 2021 Oct 5;6(41):27351-27357.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.