

PROTAC IRAK4 degrader-3 Biochemical

PROTAC IRAK4 degrader-3

CATALOG #
AAA228355


PRODUCT SYNONYMS

PROTAC IRAK4 degrader-3; PROTAC IRAK4 degrader-3 biochemical


PRODUCT SPECIFICATIONS

Purity/Purification	98%
CAS	2374122-43-5
Solubility	<1mg/ml refers to the product slightly soluble or insoluble
In vitro	PROTAC IRAK4 degrader-3 (0.01~10 uM; 2 hours; PBMCs) mediated degradation is occurring in a proteasome dependent manner[1]. PROTAC IRAK4 degrader-3 (0.01~100 uM; 18 hours; PBMCs) is capable of completely blocking IL-6 secretion[1]. Western Blot Analysis[1]Cell Line: PBMCs Concentration: 0.01~10 uM Incubation Time: 2 hours. Result: IRAK4 PROTAC-mediated degradation was occurring in a proteasome dependent manner.
Preparation and Storage	Powder: -20 degree C for 3 years In solvent: -80 degree C for 2 years

IMAGES



Structure
Dibromoacetaldehyde, CAS 3039-13-2



Structure
Dibromoacetaldehyde, CAS 3039-13-2

ADDITIONAL INFORMATION

Related Product Information for PROTAC IRAK4 degrader-3 biochemical	PROTAC IRAK4 degrader-3 is a PROTAC-induced IRAK4 degrader based on von Hippel-Lindau.
References	1. Jeong CH, et al. Occurrence and Comparative Toxicity of Haloacetaldehyde Disinfection Byproducts in Drinking Water. Environ Sci Technol. 2015;49(23):13749-13759.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.