

Human High Mobility Group Protein 1 (HMG1) ELISA Kit | HMG1 elisa kit

Human High Mobility Group Protein 1 (HMG1) ELISA Kit

CATALOG #
AAA22927

REACTIVITY
Human


PRODUCT SYNONYMS

High Mobility Group Protein 1 (HMG1); Human High Mobility Group Protein 1 (HMG1) ELISA Kit; HMGB1; HMG3; SBP1; Sulfoglucuronyl Carbohydrate Binding Protein; Amphoterin; High Mobility Group Box 1 Protein; HMG1 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of HMG1. No significant cross-reactivity or interference between HMG1 and analogues was observed.
Sequence Length	215
Samples	Human serum, plasma and other biological fluids
Detection Range	62.5-4,000pg/mL
Sensitivity	< 28.3pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level HMG1 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level HMG1 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for HMG1 elisa kit	Principle of the Assay: The microplate provided in this kit has been pre-coated with an antibody specific to HMG1. Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to HMG1. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain HMG1, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of HMG1 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	4504425
NCBI GeneID	3146
NCBI Accession #	NP_002119.1
NCBI GenBank Nucleotide #	NM_002128.5
UniProt Accession #	P09429; Q14321; Q5T7C3; Q6IBE1; A5D8W9
Molecular Weight	24,894 Da
NCBI Official Full Name	high mobility group protein B1
NCBI Official Synonym Full Names	high mobility group box 1
NCBI Official Symbol	HMGB1
NCBI Official Synonym Symbols	HMG1; HMG3; HMG-1; SBP-1
NCBI Protein Information	high mobility group protein B1
UniProt Protein Name	High mobility group protein B1
UniProt Gene Name	HMGB1
UniProt Synonym Gene Names	HMG1; HMG-1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.