

Rabbit anti-Human Membrane cofactor protein Polyclonal Antibody | anti-CD46 antibody

Rabbit anti-human Membrane cofactor protein polyclonal Antibody(CD46)

CATALOG # AAA229347	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

Membrane cofactor protein; Polyclonal Antibody; Rabbit anti-human Membrane cofactor protein polyclonal Antibody(CD46); TLX; Trophoblast leukocyte common antigen; CD46; MCP; MIC10; anti-CD46 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Caprylic Acid Ammonium Sulfate Precipitation purified
Form/Format	Liquid
Sequence Length	392
Applicable Applications for anti-CD46 antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Recombinant human Membrane cofactor protein
Conjugation	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300; Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-7056 / sc-9098
Preparation and Storage	Shipped at 4 degree C. Upon delivery, aliquot and store at -20 degree C or -80 degree C.

IMAGES



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human placenta using AAA229347 at dilution 1:100



WB (Western Blot)

Western blotAll lanes: Membrane cofactor protein antibody at 2ug/mlLane 1:K562 whole cell lysateLane 2:Hela whole cell lysateSecondaryGoat polyclonal to rabbit at 1/10000 dilutionPredicted band size: 44,45,43,41,42,40,37kDaObserved band size: 41 kDa

ADDITIONAL INFORMATION

Related Product Information for anti-CD46 antibody	Acts as a cofactor for complement factor I, a serine protease which protects autologous cells against complement-mediated injury by cleaving C3b and C4b deposited on host tissue. May be involved in the fusion of the spermatozoa with the oocyte during fertilization. Also acts as a costimulatory factor for T-cells which induces the differentiation of CD4+ into T-regulatory 1 cells. T-regulatory 1 cells suppress immune responses by secreting interleukin-10, and therefore are thought to prevent autoimmunity. A number of viral and bacterial pathogens seem to exploit this property and directly induce an immunosuppressive phenotype in T-cells by binding to CD46.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	27502402
NCBI GeneID	4179
NCBI Accession #	NP_002380.3
NCBI GenBank Nucleotide #	NM_002389.4
UniProt Accession #	P15529; Q15429; Q53GV9; Q5HY94; Q5VWS6; Q5VWS7; Q5VWS8; Q5VWS9; Q5VWT0; A0T1T0; A0T1T1; A0T1T2
Molecular Weight	36,826 Da
NCBI Official Full Name	membrane cofactor protein isoform 1
NCBI Official Synonym Full Names	CD46 molecule
NCBI Official Symbol	CD46
NCBI Official Synonym Symbols	MCP; TLX; AHUS2; MIC10; TRA2.10
NCBI Protein Information	membrane cofactor protein
UniProt Protein Name	Membrane cofactor protein
UniProt Gene Name	CD46
UniProt Synonym Gene Names	MCP; MIC10
UniProt Entry Name	MCP_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/membrane-cofactor-protein-antibody-immunohistochemistry-western-blot-elisa/229347>