

Rabbit anti-Human Tumor necrosis factor Polyclonal Antibody | anti-TNF antibody

Rabbit Anti-Human Tumor necrosis factor PolyClonal Antibody

CATALOG # AAA229601	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

Tumor necrosis factor; Polyclonal Antibody; Rabbit Anti-Human Tumor necrosis factor PolyClonal Antibody; Cachectin TNF-alpha Tumor necrosis factor ligand superfamily member 2 TNF TNFA; TNFSF2; anti-TNF antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Caprylic Acid Ammonium Sulfate Precipitation purified
Form/Format	Liquid
Sequence Length	233
Applicable Applications for anti-TNF antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Recombinant human Tumor necrosis factor protein
Conjugation	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300; Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-1351 / sc-1348 / sc-1350 / sc-8301
Preparation and Storage	Shipped at 4 degree C. Upon delivery, aliquot and store at -20 degree C or -80 degree C.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human kidney tissue using AAA229601 at dilution of 1:100




IHC (Immunohistochemisry)

Immunohistochemistry of paraffin-embedded human colon cancer using AAA229601 at dilution of 1:100



WB (Western Blot)

Western Blot Positive WB detected in: hepG2 cell, mouse heart, mouse liver. All lanes: TNF antibody at 4ug/ml. Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution. Predicted band size: 26 kDa. Observed band size: 26 kDa.



WB (Western Blot)

Western blot. All lanes: Tumor necrosis factor antibody at 2ug/ml + HeLa whole cell lysate. Secondary Goat polyclonal to rabbit at 1/10000 dilution. Predicted band size: 26kDa. Observed band size: 26kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-TNF antibody

Cytokine that binds to TNFRSF1A/TNFR1 and TNFRSF1B/TNFR. It is mainly secreted by macrophages and can induce cell death of certain tumor cell lines. It is potent pyrogen causing fever by direct action or by stimulation of interleukin-1 secretion and is implicated in the induction of cachexia. Under certain conditions it can stimulate cell proliferation and induce cell differentiation.

This gene encodes a multifunctional proinflammatory cytokine that belongs to the tumor necrosis factor (TNF) superfamily. This cytokine is mainly secreted by macrophages. It can bind to, and thus functions through its receptors TNFRSF1A/TNFR1 and TNFRSF1B/TNFR. This cytokine is involved in the regulation of a wide spectrum of biological processes including cell proliferation, differentiation, apoptosis, lipid metabolism, and coagulation. This cytokine has been implicated in a variety of diseases, including autoimmune diseases, insulin resistance, and cancer. Knockout studies in mice also suggested the neuroprotective function of this cytokine.

Product Categories/Family for anti-TNF antibody

Immunology

NCBI AND UNIPROT INFORMATION

NCBI GI #	25952111
NCBI GeneID	7124
NCBI Accession #	NP_000585.2
NCBI GenBank Nucleotide #	NM_000594.3
UniProt Accession #	P01375; O43647; Q9P1Q2; Q9UIV3
Molecular Weight	25,644 Da
NCBI Official Full Name	tumor necrosis factor
NCBI Official Synonym Full Names	tumor necrosis factor
NCBI Official Symbol	TNF
NCBI Official Synonym Symbols	DIF; TNFA; TNFSF2; TNLG1F; TNF-alpha
NCBI Protein Information	tumor necrosis factor
UniProt Protein Name	Tumor necrosis factor
UniProt Gene Name	TNF
UniProt Synonym Gene Names	TNFA; TNFSF2; TNF-a; NTF; ICD1; ICD2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/tumor-necrosis-factor-antibody-immunohistochemistry-western-blot-elisa/229601>