

Human Complement Receptor 2 (CR2) ELISA Kit | CR2 elisa kit

Human Complement Receptor 2 (CR2) ELISA Kit

CATALOG #
AAA22962

REACTIVITY
Human


PRODUCT SYNONYMS

Complement Receptor 2 (CR2); Human Complement Receptor 2 (CR2) ELISA Kit; CD21; C3DR; Complement Component(3d/Epstein Barr Virus)Receptor 2; Complement C3d receptor; Epstein-Barr virus receptor; EBV receptor; CR2 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of CR2. No significant cross-reactivity or interference between CR2 and analogues was observed.
Sequence Length	1029
Assay Type	Quantitative Sandwich
Samples	Human serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids
Detection Range	3.12-200ng/mL
Sensitivity	< 1.22ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level CR2 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level CR2 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CR2 elisa kit	Principle of the Assay: The microplate provided in this kit has been pre-coated with an antibody specific to CR2. Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to CR2. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain CR2, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of CR2 in the samples is then determined by comparing the O.D. of the samples to the standard curve. Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of CR2 in human serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	312433977
NCBI GeneID	289395
NCBI Accession #	NP_001099459.2
NCBI GenBank Nucleotide #	NM_001105989.2
NCBI Official Full Name	complement receptor type 2
NCBI Official Synonym Full Names	complement C3d receptor 2
NCBI Official Symbol	Cr2
NCBI Official Synonym Symbols	Cd21
NCBI Protein Information	complement receptor type 2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.