

Rabbit anti-Human Elongation factor 2 Polyclonal Antibody | anti-EEF2 antibody

Rabbit anti-human Elongation factor 2 polyclonal Antibody

CATALOG # AAA229684	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

Elongation factor 2; Polyclonal Antibody; Rabbit anti-human Elongation factor 2 polyclonal Antibody; EEF2; EF2; anti-EEF2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Caprylic Acid Ammonium Sulfate Precipitation purified
Form/Format	Liquid
Sequence Length	858
Applicable Applications for anti-EEF2 antibody	IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant human Elongation factor 2 protein
Conjugation	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300; Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-13003 / sc-13004 / sc-25634
Preparation and Storage	Shipped at 4 degree C. Upon delivery, aliquot and store at -20 degree C or -80 degree C.

IMAGES



IF (Immunofluorescence)

Immunofluorescent analysis of HepG2 cells using AAA229684 at a dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human kidney using AAA229684 at dilution of 1:50

ADDITIONAL INFORMATION

Related Product Information for anti-EEF2 antibody	Catalyzes the GTP-dependent ribosomal translocation step during translation elongation. During this step, the ribosome changes from the pre-translocational (PRE) to the post-translocational (POST) state as the newly formed A-site-bound peptidyl-tRNA and P-site-bound deacylated tRNA move to the P and E sites, respectively. Catalyzes the coordinated movement of the two tRNA molecules, the mRNA and conformational changes in the ribosome. This gene encodes a member of the GTP-binding translation elongation factor family. This protein is an essential factor for protein synthesis. It promotes the GTP-dependent translocation of the nascent protein chain from the A-site to the P-site of the ribosome. This protein is completely inactivated by EF-2 kinase phosphorylation.
Product Categories/Family for anti-EEF2 antibody	Epigenetics and Nuclear Signaling

NCBI AND UNIPROT INFORMATION

NCBI GI #	4503483
NCBI GeneID	1938
NCBI Accession #	NP_001952.1
NCBI GenBank Nucleotide #	NM_001961.3
UniProt Accession #	P13639; Q58J86; B2RMP5; D6W618
Molecular Weight	95,338 Da
NCBI Official Full Name	elongation factor 2
NCBI Official Synonym Full Names	eukaryotic translation elongation factor 2
NCBI Official Symbol	EEF2
NCBI Official Synonym Symbols	EF2; EF-2; EEF-2; SCA26
UniProt Protein Name	Elongation factor 2
UniProt Gene Name	EEF2
UniProt Synonym Gene Names	EF2; EF-2
UniProt Entry Name	EF2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/elongation-factor-2-antibody-immunohistochemistry-elisa/229684>