

Rabbit anti-Human S1PR3 Polyclonal Antibody | anti-S1PR3 antibody

S1PR3 Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA229729	Rabbit	Human	Immunofluorescence, Western Blot, ELISA

PRODUCT SYNONYMS


S1PR3; Polyclonal Antibody; S1PR3 Antibody; Sphingosine 1-phosphate receptor 3; S1P receptor 3; S1P3; Endothelial differentiation G-protein coupled receptor 3; Sphingosine 1-phosphate receptor Edg-3; S1P receptor Edg-3; EDG3; anti-S1PR3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	378
Applicable Applications for anti-S1PR3 antibody	IF (Immunofluorescence), WB (Western Blot), ELISA
Immunogen	Recombinant human Sphingosine 1-phosphate receptor 3 protein
Conjugation	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-22214 / sc-22210 / sc-22211 / sc-16076 / sc-30024
Preparation and Storage	Shipped at 4 degree C. Upon delivery, aliquot and store at -20 degree C or -80 degree C.



IHC (Immunohistochemistry)
Immunofluorescent analysis of MCF-7 cells using AAA229729 at a dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)



IHC (Immunohistochemistry)
IHC image of AAA229729 diluted at 1:800 and staining in paraffin-embedded human heart tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohistochemistry)
IHC image of AAA229729 diluted at 1:800 and staining in paraffin-embedded human kidney tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

ADDITIONAL INFORMATION

Related Product Information for anti-S1PR3 antibody	Receptor for the lysosphingolipid sphingosine 1-phosphate (S1P). S1P is a bioactive lysophospholipid that elicits diverse physiological effect on most types of cells and tissues. When expressed in rat HTC4 hepatoma cells, is capable of mediating S1P-induced cell proliferation and suppression of apoptosis.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	38788193
NCBI GeneID	1903
NCBI Accession #	NP_005217.2
NCBI GenBank Nucleotide #	NM_005226.3
UniProt Accession #	Q99500; Q5SQD8; Q7Z5I2
Molecular Weight	42,250 Da
NCBI Official Full Name	sphingosine 1-phosphate receptor 3
NCBI Official Synonym Full Names	sphingosine-1-phosphate receptor 3

NCBI Official Symbol	S1PR3
NCBI Official Synonym Symbols	EDG3; LPB3; S1P3; EDG-3
NCBI Protein Information	sphingosine 1-phosphate receptor 3
UniProt Protein Name	Sphingosine 1-phosphate receptor 3
UniProt Gene Name	S1PR3
UniProt Synonym Gene Names	EDG3; S1P receptor 3; S1P3; S1P receptor Edg-3
UniProt Entry Name	S1PR3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/s1pr3-antibody-immunofluorescence-western-blot-elisa/229729>