

Rabbit anti-Human, mouse YWHAE Polyclonal Antibody | anti-YWHAE antibody

YWHAE Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA229746	Rabbit	Human, mouse	Immunohistochemistry, Western Blot, ELISA


PRODUCT SYNONYMS

YWHAE; Polyclonal Antibody; YWHAE Antibody; 14-3-3E; anti-YWHAE antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, mouse
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Caprylic Acid Ammonium Sulfate Precipitation purified
Form/Format	Liquid
Sequence Length	255
Applicable Applications for anti-YWHAE antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Recombinant human 14-3-3 protein epsilon protein
Conjugation	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300; Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-31961 / sc-31962 / sc-1020
Preparation and Storage	Shipped at 4 degree C. Upon delivery, aliquot and store at -20 degree C or -80 degree C.

IMAGES




IF (Immunofluorescence)

Immunofluorescence staining of HepG2 cells with AAA229746 at 1:333, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG (H+L).




IHC (Immunohistochemistry)

IHC image of AAA229746 diluted at 1:1000 and staining in paraffin-embedded human brain tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohistochemistry)

IHC image of AAA229746 diluted at 1:1000 and staining in paraffin-embedded human adrenal gland tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



WB (Western Blot)

Western Blot Positive WB detected in: A375 whole cell lysate, HeLa whole cell lysate, SH-SY5Y whole cell lysate, Jurkat whole cell lysate, NIH/3T3 whole cell lysate, Mouse brain tissue. All lanes: YWHAE antibody at 3.3ug/ml. Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution. Predicted band size: 30,27 KDa. Observed band size: 30 KDa.

ADDITIONAL INFORMATION

Related Product Information for anti-YWHAE antibody

Adapter protein implicated in the regulation of a large spectrum of both general and specialized signaling pathways. Binds to a large number of partners, usually by recognition of a phosphoserine or phosphothreonine motif. Binding generally results in the modulation of the activity of the binding partner.

NCBI AND UNIPROT INFORMATION

NCBI GI #	5803225
NCBI GeneID	7531
NCBI Accession #	NP_006752.1
NCBI GenBank Nucleotide #	NM_006761.4
UniProt Accession #	P62258; P29360; P42655; Q4VJB6; Q53XZ5; Q63631; Q7M4R4; B3KY71; D3DTH5
Molecular Weight	26,504 Da
NCBI Official Full Name	14-3-3 protein epsilon
NCBI Official Synonym Full Names	tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein epsilon
NCBI Official Symbol	YWHAE
NCBI Official Synonym Symbols	MDS; HEL2; MDCR; KCIP-1; 14-3-3E
NCBI Protein Information	14-3-3 protein epsilon
UniProt Protein Name	14-3-3 protein epsilon
UniProt Gene Name	YWHAE
UniProt Synonym Gene Names	14-3-3E
UniProt Entry Name	1433E_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/ywhae-antibody-immunohistochemistry-western-blot-elisa/229746>