

Rabbit anti-Human Cytochrome c Polyclonal Antibody | anti-CYCS antibody

Rabbit anti-Human Cytochrome c polyclonal Antibody

CATALOG # AAA229810	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

Cytochrome c; Polyclonal Antibody; Rabbit anti-Human Cytochrome c polyclonal Antibody; CYCS; CYC; anti-CYCS antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Caprylic Acid Ammonium Sulfate Precipitation purified
Form/Format	Liquid
Sequence Length	105
Applicable Applications for anti-CYCS antibody	IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant human Cytochrome c protein
Conjugation	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300; Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-8385 / sc-7159
Preparation and Storage	Shipped at 4 degree C. Upon delivery, aliquot and store at -20 degree C or -80 degree C.

IMAGES




IHC (Immunohistochemistry)

IHC image of AAA229810 diluted at 1:200 and staining in paraffin-embedded human colon cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohistochemistry)

IHC image of AAA229810 diluted at 1:200 and staining in paraffin-embedded human kidney tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



WB (Western Blot)
 Western BlotPositive WB detected in: Rat kidney tissue,Rat heart tissue,Mouse heart tissueAll
 lanes: CYCS antibody at 2ug/mlSecondaryGoat polyclonal to rabbit IgG at 1/50000
 dilutionPredicted band size: 12 KDaObserved band size: 12 KDa

ADDITIONAL INFORMATION

Related Product Information for anti-CYCS antibody	Electron carrier protein. The oxidized form of the cytochrome c heme group can accept an electron from the heme group of the cytochrome c1 subunit of cytochrome reductase. Cytochrome c then transfers this electron to the cytochrome oxidase complex, the final protein carrier in the mitochondrial electron-transport chain.Plays a role in apoptosis. Suppression of the anti-apoptotic members or activation of the pro-apoptotic members of the Bcl-2 family leads to altered mitochondrial membrane permeability resulting in release of cytochrome c into the cytosol. Binding of cytochrome c to Apaf-1 triggers the activation of caspase-9, which then accelerates apoptosis by activating other caspases. This gene encodes cytochrome c, a component of the electron transport chain in mitochondria. The heme group of cytochrome c accepts electrons from the b-c1 complex and transfers electrons to the cytochrome oxidase complex. Cytochrome c is also involved in initiation of apoptosis. Upon release of cytochrome c to the cytoplasm, the protein binds apoptotic protease activating factor which activates the apoptotic initiator procaspase 9. Many cytochrome c pseudogenes exist, scattered throughout the human genome.
Product Categories/Family for anti-CYCS antibody	Apoptosis

NCBI AND UNIPROT INFORMATION

NCBI GI #	11128019
NCBI GeneID	54205
NCBI Accession #	NP_061820.1
NCBI GenBank Nucleotide #	NM_018947.5
UniProt Accession #	P99999; P00001; Q6NUR2; Q6NX69; Q96BV4; A4D166; B2R411
Molecular Weight	11,749 Da
NCBI Official Full Name	cytochrome c
NCBI Official Synonym Full Names	cytochrome c, somatic
NCBI Official Symbol	CYCS
NCBI Official Synonym Symbols	CYC; HCS; THC4
NCBI Protein Information	cytochrome c
UniProt Protein Name	Cytochrome c
UniProt Gene Name	CYCS

UniProt Synonym Gene Names	CYC
UniProt Entry Name	CYC_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cytochrome-c-antibody-immunohistochemistry-elisa/229810>