

Rabbit anti-Human HLA class II histocompatibility antigen, DP beta 1 chain Polyclonal Antibody | anti-HLA-DPB1 antibody

Rabbit anti-human HLA class II histocompatibility antigen, DP beta 1 chain polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA229828	Rabbit	Human	Immunohistochemistry, Western Blot, ELISA


PRODUCT SYNONYMS

HLA class II histocompatibility antigen; DP beta 1 chain; Polyclonal Antibody; Rabbit anti-human HLA class II histocompatibility antigen; DP beta 1 chain polyclonal Antibody; DP(W4) beta chain; MHC class II antigen DPB1; HLA-DPB1; HLA-DP1B; anti-HLA-DPB1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Caprylic Acid Ammonium Sulfate Precipitation purified
Form/Format	Liquid
Sequence Length	258
Applicable Applications for anti-HLA-DPB1 antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Recombinant human HLA class II histocompatibility antigen, DP beta 1 chain protein
Conjugation	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300; Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Preparation and Storage	Shipped at 4 degree C. Upon delivery, aliquot and store at -20 degree C or -80 degree C.

IMAGES



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human tonsil using AAA229828 at dilution of 1:50



WB (Western Blot)

All lanes: HLA class II histocompatibility antigen, DP beta 1 chain antibody at 2ug/mL
Lane 1: sw1990 whole cell lysate
Lane 2: HGC-27 whole cell lysate
Secondary: Goat polyclonal to Rabbit IgG at 1/10000 dilution
Predicted band size: 26kDa
Observed band size: 26kDa

ADDITIONAL INFORMATION

Related Product Information for anti-HLA-DPB1 antibody	Binds peptides derived from antigens that access the endocytic route of antigen presenting cells (APC) and presents them on the cell surface for recognition by the CD4 T-cells. The peptide binding cleft accommodates peptides of 10-30 residues. The peptides presented by MHC class II molecules are generated mostly by degradation of proteins that access the endocytic route, where they are processed by lysosomal proteases and other hydrolases. Exogenous antigens that have been endocytosed by the APC are thus readily available for presentation via MHC II molecules, and for this reason this antigen presentation pathway is usually referred to as exogenous. As membrane proteins on their way to degradation in lysosomes as part of their normal turn-over are also contained in the endosomal/lysosomal compartments, exogenous antigens must compete with those derived from endogenous components. Autophagy is also a source of endogenous peptides, autophagosomes constitutively fuse with MHC class II loading compartments. In addition to APCs, other cells of the gastrointestinal tract, such as epithelial cells, express MHC class II molecules and CD74 and act as APCs, which is an unusual trait of the GI tract. To produce a MHC class II molecule that presents an antigen, three MHC class II molecules (heterodimers of an alpha and a beta chain) associate with a CD74 trimer in the ER to form a heterononamer. Soon after the entry of this complex into the endosomal/lysosomal system where antigen processing occurs, CD74 undergoes a sequential degradation by various proteases, including CTSS and CTSL, leaving a small fragment termed CLIP (class-II-associated invariant chain peptide). The removal of CLIP is facilitated by HLA-DM via direct binding to the alpha-beta-CLIP complex so that CLIP is released. HLA-DM stabilizes MHC class II molecules until primary high affinity antigenic peptides are bound. The MHC II molecule bound to a peptide is then transported to the cell membrane surface. In B-cells, the interaction between HLA-DM and MHC class II molecules is regulated by HLA-DO. Primary dendritic cells (DCs) also to express HLA-DO. Lysosomal miroenvironment has been implicated in the regulation of antigen loading into MHC II molecules, increased acidification produces increased proteolysis and efficient peptide loading. HLA-DPB belongs to the HLA class II beta chain paralogues. This class II molecule is a heterodimer consisting of an alpha (DPA) and a beta chain (DPB), both anchored in the membrane. It plays a central role in the immune system by presenting peptides derived from extracellular proteins. Class II molecules are expressed in antigen presenting cells (APC: B lymphocytes, dendritic cells, macrophages). The beta chain is approximately 26-28 kDa and its gene contains 6 exons. Exon one encodes the leader peptide, exons 2 and 3 encode the two extracellular domains, exon 4 encodes the transmembrane domain and exon 5 encodes the cytoplasmic tail. Within the DP molecule both the alpha chain and the beta chain contain the polymorphisms specifying the peptide binding specificities, resulting in up to 4 different molecules.
Product Categories/Family for anti-HLA-DPB1 antibody	Immunology

NCBI AND UNIPROT INFORMATION

NCBI GI #	24797076
NCBI GeneID	3115
NCBI Accession #	NP_002112.3
NCBI GenBank Nucleotide #	NM_002121.5
UniProt Accession #	P04440; A0PFJ7; A5I886; A8YPB3; B5U8B4; B7VF80; B7VF87; B8ZX68; B8ZYT0; B9W5S8; B9W6F7; B9W6F9
Molecular Weight	29,159 Da
NCBI Official Full Name	HLA class II histocompatibility antigen, DP beta 1 chain
NCBI Official Synonym Full Names	major histocompatibility complex, class II, DP beta 1
NCBI Official Symbol	HLA-DPB1
NCBI Official Synonym Symbols	DPB1; HLA-DP; HLA-DPB; HLA-DP1B
NCBI Protein Information	HLA class II histocompatibility antigen, DP beta 1 chain
UniProt Protein Name	HLA class II histocompatibility antigen, DP beta 1 chain
UniProt Gene Name	HLA-DPB1
UniProt Synonym Gene Names	HLA-DP1B

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/hla-class-ii-histocompatibility-antigen-dp-beta-1-chain-antibody-immunohistochemistry-western-blot-elisa/229828>