

Mouse Eosinophil Peroxidase (EPX) ELISA Kit | EPX elisa kit

Mouse Eosinophil Peroxidase (EPX) ELISA Kit

CATALOG #
AAA23073

REACTIVITY
Mouse


PRODUCT SYNONYMS

Eosinophil Peroxidase (EPX); Mouse Eosinophil Peroxidase (EPX) ELISA Kit; EPO; EPP; EPX-PEN; EPER; EPX elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of EPX. No significant cross-reactivity or interference between EPX and analogues was observed.
Sequence Length	715
Assay Type	Quantitative Sandwich
Samples	Mouse serum, plasma, tissue homogenates and other biological fluids
Detection Range	15.6-1,000pg/mL
Sensitivity	< 5.5pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level EPX were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level EPX were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for EPX elisa kit	Principle of the Assay: The microplate provided in this kit has been pre-coated with an antibody specific to EPX. Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to EPX. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain EPX, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of EPX in the samples is then determined by comparing the O.D. of the samples to the standard curve. Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of EPX in mouse serum, plasma, tissue homogenates and other biological fluids.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	182146
NCBI GeneID	8288
UniProt Accession #	P11678; Q4TVP3
Molecular Weight	81,040 Da
NCBI Official Full Name	eosinophil peroxidase
NCBI Official Synonym Full Names	eosinophil peroxidase
NCBI Official Symbol	EPX
NCBI Official Synonym Symbols	EPO; EPP; EPXD; EPX-PEN
NCBI Protein Information	eosinophil peroxidase
UniProt Protein Name	Eosinophil peroxidase
UniProt Gene Name	EPX
UniProt Synonym Gene Names	EPER; EPO; EPP; EPO

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.