

Rabbit anti-Human, Rat FZD3 Polyclonal Antibody | anti-FZD3 antibody

FZD3 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA230859	Rabbit	Human, Rat	Immunohistochemistry, Western Blot, ELISA

PRODUCT SYNONYMS

FZD3; Polyclonal Antibody; FZD3 Antibody; Frizzled-3; Fz-3; hFz3; anti-FZD3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen Affinity Purified
Form/Format	Liquid
Sequence Length	666
Applicable Applications for anti-FZD3 antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Species	Human
Immunogen	Recombinant human Frizzled-3 protein (23-205AA)
Conjugate	Non-conjugated
Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-68332 / sc-68333 / sc-68334 / sc-292246
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human skeletal muscle tissue AAA230859 at dilution 1:100



WB (Western Blot)

Western blotAll lanes: FZD3 antibody at 1.94ug/mlLane 1: Rat heart tissueLane 2: U251 whole cell lysateLane 3: Thp-1 whole cell lysateLane 4: 293T whole cell lysateLane 5: Hela whole cell lysateLane 6: HepG-2 whole cell lysateSecondaryGoat polyclonal to rabbit IgG at 1/10000 dilutionPredicted band size: 76 kDaObserved band size: 76 kDa

ADDITIONAL INFORMATION

Related Product Information for anti-FZD3 antibody

Receptor for Wnt proteins. Most of frizzled receptors are coupled to the beta-catenin canonical signaling pathway, which leads to the activation of disheveled proteins, inhibition of GSK-3 kinase, nuclear accumulation of beta-catenin and activation of Wnt target genes. A second signaling pathway involving PKC and calcium fluxes has been seen for some family members, but it is not yet clear if it represents a distinct pathway or if it can be integrated in the canonical pathway, as PKC seems to be required for Wnt-mediated inactivation of GSK-3 kinase. Both pathways seem to involve interactions with G-proteins. Activation by Wnt5A stimulates PKC activity via a G-protein-dependent mechanism. Involved in transduction and intercellular transmission of polarity information during tissue morphogenesis and/or in differentiated tissues. Plays a role in controlling early axon growth and guidance processes necessary for the formation of a subset of central and peripheral major fiber tracts. Required for the development of major fiber tracts in the central nervous system, including: the anterior commissure, the corpus callosum, the thalamocortical, corticothalamic and nigrostriatal tracts, the corticospinal tract, the fasciculus retroflexus, the mammillothalamic tract, the medial lemniscus, and ascending fiber tracts from the spinal cord to the brain. In the peripheral nervous system, controls axon growth in distinct populations of cranial and spinal motor neurons, including the facial branchiomotor nerve, the hypoglossal nerve, the phrenic nerve, and motor nerves innervating dorsal limbs. Involved in the migration of cranial neural crest cells. May also be implicated in the transmission of sensory information from the trunk and limbs to the brain. Controls commissural sensory axons guidance after midline crossing along the anterior-posterior axis in the developing spinal cord in a Wnt-dependent signaling pathway. Together with FZD6, is involved in the neural tube closure and plays a role in the regulation of the establishment of planar cell polarity (PCP), particularly in the orientation of asymmetric bundles of stereocilia on the apical faces of a subset of auditory and vestibular sensory cells located in the inner ear. Promotes neurogenesis by maintaining sympathetic neuroblasts within the cell cycle in a beta-catenin-dependent manner (By similarity).

NCBI AND UNIPROT INFORMATION

NCBI GI #	8393378
NCBI GeneID	7976
NCBI Accession #	NP_059108.1
NCBI GenBank Nucleotide #	NM_017412.3
UniProt Accession #	Q9NPG1; A8K615
Molecular Weight	– Da
NCBI Official Full Name	frizzled-3
NCBI Official Synonym Full Names	frizzled class receptor 3
NCBI Official Symbol	FZD3
NCBI Official Synonym Symbols	Fz-3
NCBI Protein Information	frizzled-3
UniProt Protein Name	Frizzled-3
UniProt Gene Name	FZD3
UniProt Synonym Gene Names	Fz-3; hFz3
UniProt Entry Name	FZD3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/fzd3-antibody-immunohistochemistry-western-blot-elisa/230859>