

Rabbit anti-Human FCER1A Polyclonal Antibody | anti-FCER1A antibody

FCER1A Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA231156	Rabbit	Human	Immunofluorescence, Immunohistochemistry, Western Blot, ELISA

PRODUCT SYNONYMS


FCER1A; Polyclonal Antibody; FCER1A Antibody; High affinity immunoglobulin epsilon receptor subunit alpha; Fc-epsilon RI-alpha; FcERI; IgE Fc receptor subunit alpha; FCE1A; anti-FCER1A antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	257
Applicable Applications for anti-FCER1A antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot), ELISA
Species	Human
Immunogen	Recombinant human High affinity immunoglobulin epsilon receptor subunit alpha protein (26-205AA)
Conjugate	Non-conjugated
Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-33480 / sc-33482 / sc-68942
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.




WB (Western Blot)
Western BlotPositive WB detected in:Mouse heart tissue,Mouse brain tissueAll lanes: FCER1A antibody at 3ug/mlSecondaryGoat polyclonal to rabbit IgG at 1/50000 dilutionPredicted band size: 30 kDaObserved band size: 30 kDa



IHC (Immunohistochemisry)
Immunofluorescent analysis of Hela cells using AAA231156 at a dilution of 1:100 and Alexa Fluor 488-conugated AffiniPure Goat Anti-Rabbit IgG(H+L)



IHC (Immunohiostchemistry)
Immunohistochemistry of paraffin-embedded human skin tissue using AAA231156 at dilution of 1:100



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded human prostate cancer using AAA231156 at dilution of 1:100

ADDITIONAL INFORMATION

Related Product Information for anti-FCER1A antibody	Binds to the Fc region of immunoglobulins epsilon. High affinity receptor. Responsible for initiating the allergic response. Binding of allergen to receptor-bound IgE leads to cell activation and the release of mediators (such as histamine) responsible for the manifestations of allergy. The same receptor also induces the secretion of important lymphokines.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4503675
NCBI GeneID	2205
NCBI Accession #	NP_001992.1
NCBI GenBank Nucleotide #	NM_002001.3
UniProt Accession #	P12319
Molecular Weight	29,596 Da
NCBI Official Full Name	high affinity immunoglobulin epsilon receptor subunit alpha
NCBI Official Synonym Full Names	Fc fragment of IgE receptor Ia
NCBI Official Symbol	FCER1A

NCBI Official Synonym Symbols	FCE1A; FcERI
NCBI Protein Information	high affinity immunoglobulin epsilon receptor subunit alpha
UniProt Protein Name	High affinity immunoglobulin epsilon receptor subunit alpha
UniProt Gene Name	FCER1A
UniProt Synonym Gene Names	FCE1A; FcERI
UniProt Entry Name	FCERA_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/fcer1a-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/231156>