

Rabbit anti-Human CARM1 Polyclonal Antibody | anti-CARM1 antibody

CARM1 Antibody, HRP conjugated

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA231184	Rabbit	Human	ELISA

PRODUCT SYNONYMS

CARM1; Polyclonal Antibody; CARM1 Antibody; HRP conjugated; Histone-arginine methyltransferase CARM1; Coactivator-associated arginine methyltransferase 1; Protein arginine N-methyltransferase 4; PRMT4; anti-CARM1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	608
Applicable Applications for anti-CARM1 antibody	ELISA
Species	Human
Immunogen	Recombinant human Histone-arginine methyltransferase CARM1 protein
Conjugate	HRP
Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-33176
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.

ADDITIONAL INFORMATION

Related Product Information for anti-CARM1 antibody	Methylates (mono- and asymmetric dimethylation) the guanidino nitrogens of arginyl residues in several proteins involved in DNA packaging, transcription regulation, pre-mRNA splicing, and mRNA stability. Recruited to promoters upon gene activation together with histone acetyltransferases from EP300/P300 and p160 families, methylates histone H3 at 'Arg-17' (H3R17me), forming mainly asymmetric dimethylarginine (H3R17me2a), leading to activate transcription via chromatin remodeling. During nuclear hormone receptor activation and TCF7L2/TCF4 activation, acts synergically with EP300/P300 and either one of the p160 histone acetyltransferases NCOA1/SRC1, NCOA2/GRIP1 and NCOA3/ACTR or CTNNB1/beta-catenin to activate transcription. During myogenic transcriptional activation, acts together with NCOA3/ACTR as a coactivator for MEF2C. During monocyte inflammatory stimulation, acts together with EP300/P300 as a coactivator for NF-kappa-B. Acts as coactivator for PPARG, promotes adipocyte differentiation and the accumulation of brown fat tissue. Plays a role in the regulation of pre-mRNA alternative splicing by methylation of splicing factors. Also seems to be involved in p53/TP53 transcriptional activation. Methylates EP300/P300, both at 'Arg-2142', which may loosen its interaction with NCOA2/GRIP1, and at 'Arg-580' and 'Arg-604' in the KIX domain, which impairs its interaction with CREB and inhibits CREB-dependent transcriptional activation. Also methylates arginine residues in RNA-binding proteins PABPC1, ELAVL1 and ELAV4, which may affect their mRNA-stabilizing properties and the half-life of their target mRNAs.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	40288288
NCBI GeneID	10498
NCBI Accession #	NP_954592.1
NCBI GenBank Nucleotide #	NM_199141.1
UniProt Accession #	Q86X55; A6NN38
Molecular Weight	41,885 Da
NCBI Official Full Name	histone-arginine methyltransferase CARM1
NCBI Official Synonym Full Names	coactivator associated arginine methyltransferase 1
NCBI Official Symbol	CARM1
NCBI Official Synonym Symbols	PRMT4
NCBI Protein Information	histone-arginine methyltransferase CARM1
UniProt Protein Name	Histone-arginine methyltransferase CARM1
UniProt Gene Name	CARM1
UniProt Synonym Gene Names	PRMT4
UniProt Entry Name	CARM1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.