

Rabbit anti-Human CD276 Polyclonal Antibody | anti-CD276 antibody

CD276 Antibody

CATALOG # AAA231345	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

CD276; Polyclonal Antibody; CD276 Antibody; CD276 antigen; 4lg-B7-H3; B7 homolog 3; B7-H3; Costimulatory molecule; B7H3; PSEC0249; UNQ309/PRO352; anti-CD276 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen Affinity Purified
Form/Format	Liquid
Sequence Length	534
Applicable Applications for anti-CD276 antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Species	Human
Immunogen	Recombinant human CD276 antigen protein (29-200AA)
Conjugate	Non-conjugated
Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-34569 / sc-66814
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IHC (Immunohiostchemistry)
Immunohistochemistry of paraffin-embedded human prostate cancer AAA231345 at dilution 1:100



WB (Western Blot)
Western blotAll lanes: CD276 antibody at 0.48ug/mlLane 1: U251 whole cell lysateLane 2: Jurkat whole cell lysateLane 3: HepG-2 whole cell lysateLane 4: 293T whole cell lysateSecondaryGoat polyclonal to rabbit IgG at 1/10000 dilutionPredicted band size: 58,34,53 kDaObserved band size: 57 kDa

ADDITIONAL INFORMATION

Related Product Information for anti-CD276 antibody	May participate in the regulation of T-cell-mediated immune response. May play a protective role in tumor cells by inhibiting natural-killer mediated cell lysis as well as a role of marker for detection of neuroblastoma cells. May be involved in the development of acute and chronic transplant rejection and in the regulation of lymphocytic activity at mucosal surfaces. Could also play a key role in providing the placenta and fetus with a suitable immunological environment throughout pregnancy. Both isoform 1 and isoform 2 appear to be redundant in their ability to modulate CD4 T-cell responses. Isoform 2 is shown to enhance the induction of cytotoxic T-cells and selectively stimulates interferon gamma production in the presence of T-cell receptor signaling.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	67188443
NCBI GeneID	80381
NCBI Accession #	NP_001019907.1
NCBI GenBank Nucleotide #	NM_001024736.1
UniProt Accession #	Q5ZPR3; Q6P5Y4; Q6UXI2; Q8NBI8; Q8NC34; Q8NCB6; Q9BXR1
Molecular Weight	57,165 Da
NCBI Official Full Name	CD276 antigen isoform a
NCBI Official Synonym Full Names	CD276 molecule
NCBI Official Symbol	CD276
NCBI Official Synonym Symbols	B7H3; B7-H3; B7RP-2; 4Ig-B7-H3
NCBI Protein Information	CD276 antigen
UniProt Protein Name	CD276 antigen
UniProt Gene Name	CD276
UniProt Synonym Gene Names	B7H3; B7-H3
UniProt Entry Name	CD276_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/cd276-antibody-immunohistochemistry-western-blot-elisa/231345>