

Rabbit anti-Human STAT3 Polyclonal Antibody | anti-STAT3 antibody

STAT3 Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA231554	Rabbit	Human	Immunofluorescence, Immunohistochemistry, ELISA

PRODUCT SYNONYMS

STAT3; Polyclonal Antibody; STAT3 Antibody; Signal transducer and activator of transcription 3; Acute-phase response factor; anti-STAT3 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	769
Applicable Applications for anti-STAT3 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Species	Human
Immunogen	Recombinant human Signal transducer and activator of transcription 3 protein (50-240AA)
Conjugate	Non-conjugated
Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-483 / sc-482 / sc-7179
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.




IP (Immunoprecipitation)

Immunoprecipitating STAT3 in Hela whole cell lysate Lane 1: Rabbit control IgG (1ug) instead of AAA231554 in Hela whole cell lysate. For western blotting, a HRP-conjugated Protein G antibody was used as the secondary antibody (1/2000) Lane 2: AAA231554 (6ug)+ Hela whole cell lysate (500ug) Lane 3: Hela whole cell lysate (10ug)



IF (Immunofluorescence)

Immunofluorescent analysis of HepG-2 cells using AAA231554 at a dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human liver tissue using AAA231554 at dilution 1:100

ADDITIONAL INFORMATION

Related Product Information for anti-STAT3 antibody

Signal transducer and transcription activator that mediates cellular responses to interleukins, KITLG/SCF, LEP and other growth factors. Once activated, recruits coactivators, such as NCOA1 or MED1, to the promoter region of the target gene (PubMed:17344214). May mediate cellular responses to activated FGFR1, FGFR2, FGFR3 and FGFR4. Binds to the interleukin-6 (IL-6)-responsive elements identified in the promoters of various acute-phase protein genes. Activated by IL31 through IL31RA. Involved in cell cycle regulation by inducing the expression of key genes for the progression from G1 to S phase, such as CCND1 (PubMed:17344214). Mediates the effects of LEP on melanocortin production, body energy homeostasis and lactation (By similarity). May play an apoptotic role by transactivating BIRC5 expression under LEP activation (PubMed:18242580). Cytoplasmic STAT3 represses macroautophagy by inhibiting EIF2AK2/PKR activity.

NCBI AND UNIPROT INFORMATION

NCBI GI #	21618338
NCBI GeneID	6774
NCBI Accession #	NP_003141.2
NCBI GenBank Nucleotide #	NM_003150.3
UniProt Accession #	P40763; O14916; Q9BW54; A8K7B8; K7ENL3
Molecular Weight	83,126 Da
NCBI Official Full Name	signal transducer and activator of transcription 3 isoform 2

NCBI Official Synonym Full Names	signal transducer and activator of transcription 3
NCBI Official Symbol	STAT3
NCBI Official Synonym Symbols	APRF; HIES; ADMIO; ADMIO1
NCBI Protein Information	signal transducer and activator of transcription 3
UniProt Protein Name	Signal transducer and activator of transcription 3
UniProt Gene Name	STAT3
UniProt Entry Name	STAT3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/stat3-antibody-immunofluorescence-immunohistochemistry-elisa/231554>