

Rabbit anti-Human TAF9 Polyclonal Antibody | anti-TAF9 antibody

TAF9 Antibody

CATALOG # AAA231871	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

TAF9; Polyclonal Antibody; TAF9 Antibody; Transcription initiation factor TFIID subunit 9; RNA polymerase II TBP-associated factor subunit G; STAF31/32; Transcription initiation factor TFIID 31 kDa subunit; TAFII-31; TAFII31; Transcription initiation factor TFIID 32 kDa subunit; TAFII-32; TAFII32; TAF2G; anti-TAF9 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen Affinity Purified
Form/Format	Liquid
Sequence Length	264
Applicable Applications for anti-TAF9 antibody	IHC (Immunohistochemistry), ELISA
Species	Human
Immunogen	Recombinant human Transcription initiation factor TFIID subunit 9 protein (1-172AA)
Conjugate	Non-conjugated
Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human skin tissue using AAA231871 at dilution 1:100



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human thyroid tissue using AAA231871 at dilution 1:100



WB (Western Blot)

Western BlotPositive WB detected in:MCF-7 whole cell lysate,293T whole cell lysate,HeLa whole cell lysate,K562 whole cell lysate,A431 whole cell lysateAll lanes: AK6 antibody at 0.59ug/mlSecondaryGoat polyclonal to rabbit IgG at 1/50000 dilutionPredicted band size: 21,20kDaObserved band size: 20 kDa

ADDITIONAL INFORMATION

Related Product Information for anti-TAF9 antibody

Essential for cell viability. TAF9 and TAF9B are involved in transcriptional activation as well as repression of distinct but overlapping sets of genes. May have a role in gene regulation associated with apoptosis. TAFs are components of the transcription factor IID (TFIID) complex, the TBP-free TAFII complex (TFTC), the PCAF histone acetylase complex and the STAGA transcription coactivator-HAT complex. TFIID or TFTC are essential for the regulation of RNA polymerase II-mediated transcription.

NCBI AND UNIPROT INFORMATION

NCBI GI #	62865614
NCBI GeneID	6880
NCBI Accession #	NP_001015892.1
NCBI GenBank Nucleotide #	NM_001015892.1
UniProt Accession #	Q16594; Q5U0D1; Q9BTS1; D3DWA3
Molecular Weight	28,974 Da
NCBI Official Full Name	transcription initiation factor TFIID subunit 9
NCBI Official Synonym Full Names	TATA-box binding protein associated factor 9
NCBI Official Symbol	TAF9
NCBI Official Synonym Symbols	TAF2G; TAFII31; TAFII32; MGC:5067; TAFII-31; TAFII-32; TAFIID32; STAF31/32
NCBI Protein Information	transcription initiation factor TFIID subunit 9
UniProt Protein Name	Transcription initiation factor TFIID subunit 9
UniProt Gene Name	TAF9
UniProt Synonym Gene Names	TAF2G; TAFII31; TAFII-31; TAFII31; TAFII-32; TAFII32
UniProt Entry Name	TAF9_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/taf9-antibody-immunohistochemistry-elisa/231871>