

Rabbit anti-Human GRIA4 Polyclonal Antibody | anti-GRIA4 antibody

GRIA4 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA232511	Rabbit	Human	Immunofluorescence, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

GRIA4; Polyclonal Antibody; GRIA4 Antibody; Glutamate receptor 4GluR-4GluR4; AMPA-selective glutamate receptor 4; GluR-D; Glutamate receptor ionotropic; AMPA 4; GluA4; GLUR4; anti-GRIA4 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	902
Applicable Applications for anti-GRIA4 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant human Glutamate receptor 4 protein (733-883AA)
Preservative	0.03% Proclin 300
Constituents	50% Glycerol, 0.01M PBS, PH 7.4
Conjugation	Non-conjugated
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-31394 / sc-7614
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES


IF (Immunofluorescence)

Immunofluorescent analysis of HepG2 cells using AAA232511 at a dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)


IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human brain tissue using AAA232511 at dilution of 1:100

ADDITIONAL INFORMATION

Related Product Information for anti-GRIA4 antibody	Receptor for glutamate that functions as ligand-gated ion channel in the central nervous system and plays an important role in excitatory synaptic transmission. L-glutamate acts as an excitatory neurotransmitter at many synapses in the central nervous system. Binding of the excitatory neurotransmitter L-glutamate induces a conformation change, leading to the opening of the cation channel, and thereby converts the chemical signal to an electrical impulse. The receptor then desensitizes rapidly and enters a transient inactive state, characterized by the presence of bound agonist. In the presence of CACNG4 or CACNG7 or CACNG8, shows resensitization which is characterized by a delayed accumulation of current flux upon continued application of glutamate.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	164419734
NCBI GeneID	2893
NCBI Accession #	NP_000820.3
NCBI GenBank Nucleotide #	NM_000829.3
UniProt Accession #	P48058; Q86XE8
Molecular Weight	49,146 Da
NCBI Official Full Name	glutamate receptor 4 isoform 1
NCBI Official Synonym Full Names	glutamate ionotropic receptor AMPA type subunit 4
NCBI Official Symbol	GRIA4
NCBI Official Synonym Symbols	GLUR4; GLURD; GluA4; GLUR4C
NCBI Protein Information	glutamate receptor 4
UniProt Protein Name	Glutamate receptor 4
UniProt Gene Name	GRIA4
UniProt Synonym Gene Names	GLUR4; GluR-4; GluR4; GluA4
UniProt Entry Name	GRIA4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/gria4-antibody-immunofluorescence-immunohistochemistry-elisa/232511>