

Rabbit anti-Human MET Polyclonal Antibody | anti-MET antibody

MET Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA232581	Rabbit	Human	Immunofluorescence, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

MET; Polyclonal Antibody; MET Antibody; Hepatocyte growth factor receptor; HGF receptor; HGF/SF receptor; Proto-oncogene c-Met; Scatter factor receptor; SF receptor; Tyrosine-protein kinase Met; anti-MET antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	1390
Applicable Applications for anti-MET antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant Human Hepatocyte growth factor receptor protein (83-209AA)
Preservative	0.03% Proclin 300
Constituents	50% Glycerol, 0.01M PBS, pH 7.4
Conjugation	Non-conjugated
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-10 / sc-161 / sc-162 / sc-34405 / sc-34407 / sc-46394 / sc-46395 / sc-8307
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




IHC (Immunohistochemistry)

Immunofluorescent analysis of U251 cells using AAA232581 at a dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human liver cancer using AAA232581 at dilution of 1:100



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded human colon cancer using AAA232581 at dilution of 1:100

ADDITIONAL INFORMATION

Related Product Information for anti-MET antibody	Receptor tyrosine kinase that transduces signals from the extracellular matrix into the cytoplasm by binding to hepatocyte growth factor/HGF ligand. Regulates many physiological processes including proliferation, scattering, morphogenesis and survival. Ligand binding at the cell surface induces autophosphorylation of MET on its intracellular domain that provides docking sites for downstream signaling molecules. Following activation by ligand, interacts with the PI3-kinase subunit PIK3R1, PLCG1, SRC, GRB2, STAT3 or the adapter GAB1. Recruitment of these downstream effectors by MET leads to the activation of several signaling cascades including the RAS-ERK, PI3 kinase-AKT, or PLCgamma-PKC. The RAS-ERK activation is associated with the morphogenetic effects while PI3K/AKT coordinates prosurvival effects. During embryonic development, MET signaling plays a role in gastrulation, development and migration of muscles and neuronal precursors, angiogenesis and kidney formation. In adults, participates in wound healing as well as organ regeneration and tissue remodeling. Promotes also differentiation and proliferation of hematopoietic cells. May regulate cortical bone osteogenesis (By similarity).
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	42741655
NCBI GeneID	4233
NCBI Accession #	NP_000236.2
NCBI GenBank Nucleotide #	NM_000245.3
UniProt Accession #	P08581; O60366; Q12875; Q9UDX7; Q9UPL8; A1L467; B5A932; E7EQ94
Molecular Weight	85,745 Da
NCBI Official Full Name	hepatocyte growth factor receptor isoform b preproprotein
NCBI Official Synonym Full Names	MET proto-oncogene, receptor tyrosine kinase
NCBI Official Symbol	MET
NCBI Official Synonym Symbols	HGFR; AUTS9; RCCP2; c-Met; DFNB97
NCBI Protein Information	hepatocyte growth factor receptor
UniProt Protein Name	Hepatocyte growth factor receptor
UniProt Gene Name	MET
UniProt Synonym Gene Names	HGF receptor; SF receptor
UniProt Entry Name	MET_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

