

Rabbit anti-Human, Mouse PAK3 Polyclonal Antibody | anti-PAK3 antibody

PAK3 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA232616	Rabbit	Human, Mouse	Immunofluorescence, Immunohistochemistry, Western Blot, ELISA

PRODUCT SYNONYMS

PAK3; Polyclonal Antibody; PAK3 Antibody; Serine/threonine-protein kinase PAK 3; Beta-PAK; Oligophrenin-3; p21-activated kinase 3; PAK-3; OPHN3; anti-PAK3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	544
Applicable Applications for anti-PAK3 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Recombinant Human Serine/threonine-protein kinase PAK 3 protein (70-190AA)
Preservative	0.03% Proclin 300
Constituents	50% Glycerol, 0.01M PBS, PH 7.4
Conjugation	Non-conjugated
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-1871
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




IHC (Immunohistochemistry)

Immunofluorescent analysis of Hela cells using AAA232616 at a dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human brain tissue using AAA232616 at dilution of 1:100



WB (Western Blot)

Western BlotPositive WB detected in:Mouse lung tissue,Mouse skeletal muscle tissueAll lanes: PAK3 antibody at 3ug/mlSecondaryGoat polyclonal to rabbit IgG at 1/50000 dilutionPredicted band size: 63,61,65 kDaObserved band size: 63 kDa

ADDITIONAL INFORMATION

Related Product Information for anti-PAK3 antibody	Serine/threonine protein kinase that plays a role in a variety of different signaling pathways including cytoskeleton regulation, cell migration, or cell cycle regulation. Plays a role in dendrite spine morphogenesis as well as synapse formation and plasticity. Acts as downstream effector of the small GTPases CDC42 and RAC1. Activation by the binding of active CDC42 and RAC1 results in a conformational change and a subsequent autophosphorylation on several serine and/or threonine residues. Phosphorylates MAPK4 and MAPK6 and activates the downstream target MAPKAPK5, a regulator of F-actin polymerization and cell migration. Additionally, phosphorylates TNNI3/troponin I to modulate calcium sensitivity and relaxation kinetics of thin myofilaments. May also be involved in early neuronal development.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	189491751
NCBI GeneID	5063
NCBI Accession #	NP_001121638.1
NCBI GenBank Nucleotide #	NM_001128166.2
UniProt Accession #	O75914; Q5JWX1; Q5JWX2; Q7Z2D6; Q7Z2E4; Q7Z3Z8; Q8WWK5; Q8WX23; A8K389; B1GX77; B1GX78; B1GX79
Molecular Weight	62,913 Da
NCBI Official Full Name	serine/threonine-protein kinase PAK 3 isoform a
NCBI Official Synonym Full Names	p21 (RAC1) activated kinase 3
NCBI Official Symbol	PAK3
NCBI Official Synonym Symbols	ARA; bPAK; MRX30; MRX47; OPHN3; PAK-3; PAK3beta; beta-PAK
NCBI Protein Information	serine/threonine-protein kinase PAK 3
UniProt Protein Name	Serine/threonine-protein kinase PAK 3
UniProt Gene Name	PAK3
UniProt Synonym Gene Names	OPHN3; PAK-3
UniProt Entry Name	PAK3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/pak3-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/232616>