

Rabbit anti-Human RAB7A Polyclonal Antibody | anti-RAB7A antibody

RAB7A Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA232647	Rabbit	Human	Immunofluorescence, Immunohistochemistry, ELISA

PRODUCT SYNONYMS

RAB7A; Polyclonal Antibody; RAB7A Antibody; Ras-related protein Rab-7a; RAB7; anti-RAB7A antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	207
Applicable Applications for anti-RAB7A antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant human Ras-related protein Rab-7a protein (110-177AA)
Preservative	0.03% Proclin 300
Constituents	50% Glycerol, 0.01M PBS, PH 7.4
Conjugation	Non-conjugated
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-11303 / sc-6563 / sc-10767
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IF (Immunofluorescence)

Immunofluorescent analysis of HepG2 cells using AAA232647 at a dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human gastric cancer tissue using AAA232647 at dilution of 1:100

ADDITIONAL INFORMATION

Related Product Information for anti-RAB7A antibody	Key regulator in endo-lysosomal trafficking. Governs early-to-late endosomal maturation, microtubule minus-end as well as plus-end directed endosomal migration and positioning, and endosome-lysosome transport through different protein-protein interaction cascades. Plays a central role, not only in endosomal traffic, but also in many other cellular and physiological events, such as growth-factor-mediated cell signaling, nutrient-transporter mediated nutrient uptake, neurotrophin transport in the axons of neurons and lipid metabolism. Also involved in regulation of some specialized endosomal membrane trafficking, such as maturation of melanosomes, pathogen-induced phagosomes (or vacuoles) and autophagosomes. Plays a role in the maturation and acidification of phagosomes that engulf pathogens, such as S.aureus and M.tuberculosis. Plays a role in the fusion of phagosomes with lysosomes. Plays important roles in microbial pathogen infection and survival, as well as in participating in the life cycle of viruses. Microbial pathogens possess survival strategies governed by RAB7A, sometimes by employing RAB7A function (e.g. Salmonella) and sometimes by excluding RAB7A function (e.g. Mycobacterium). In concert with RAC1, plays a role in regulating the formation of RBs (ruffled borders) in osteoclasts. Controls the endosomal trafficking and neurite outgrowth signaling of NTRK1/TRKA (PubMed: 11179213, PubMed: 12944476, PubMed: 14617358, PubMed: 20028791, PubMed: 21255211). Regulates the endocytic trafficking of the EGF-EGFR complex by regulating its lysosomal degradation. Involved in the ADRB2-stimulated lipolysis through lipophagy, a cytosolic lipase-independent autophagic pathway (By similarity). Required for the exosomal release of SDCBP, CD63 and syndecan (PubMed: 22660413).
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	34147513
NCBI GeneID	7879
NCBI Accession #	NP_004628.4
NCBI GenBank Nucleotide #	NM_004637.5
UniProt Accession #	P51149; Q9NWX0; Q9UPB0; A8K3V6
Molecular Weight	23,490 Da
NCBI Official Full Name	ras-related protein Rab-7a
NCBI Official Synonym Full Names	RAB7A, member RAS oncogene family
NCBI Official Symbol	RAB7A
NCBI Official Synonym Symbols	RAB7; PRO2706
NCBI Protein Information	ras-related protein Rab-7a
UniProt Protein Name	Ras-related protein Rab-7a
UniProt Gene Name	RAB7A
UniProt Synonym Gene Names	RAB7
UniProt Entry Name	RAB7A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.